FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY

formula 4216

ACTIVE INGREDIENTS:

| Nonylphenoxypoly(ethoxy) ethanol-iodine complex | 17.24% |
|---|--------|
| Phosphoric Acid | 0.60% |
| INERT INGREDIENT: | |
| Water | |

U. S. Patent Nos. 2931777, 3028299



KEEP FROM FREEZING

USDA Reg. No. 52-127

CAUTION: Keep Out Of Reach Of Children.

(See other precautions on rear panel)

manufactured by WEST CHEMICAL PRODUCTS, INC.

NET CONTENTS: 1 GALLON U.S.

42-16 West Street • Long Island City N.Y. 11101
branches in principal cities

K

FORMULA 4216 is a

DIRECTIONS FOR USE: For cleaning and dis in a single operation, use three (3) ounce five (5) gallons of water. Use a scrub brusinsure thorough cleaning. For porous surfaclean, use six (6) of ounces of FORMULA of water.

germicidal activity. In use dilution, it ex Just as long as color remains in this solut germicidal action is assured. Once colo solution should be discarded.

HOSPITAL USE: Use three (3) ounces of F(gallons of water (75 PPM titratable iodin infection, isolation and terminal disinference of rubber sheets and rubber glove (5 minutes contact recommended.)

NLY

.24%

.60%

.16%

.0%

52-127

INC.

Y. 11101

al cities

formula 4216

FORMULA 4216 is a general purpose "Tamed Iodine" detergent germicide applicable for a variety of uses

DIRECTIONS FOR USE: For cleaning and disinfecting most surfaces in a single operation, use three (3) ounces of FORMULA 4216 to five (5) gallons of water. Use a scrub brush or suitable device to insure thorough cleaning. For porous surfaces or those difficult to clean, use six (6) of ounces of FORMULA 4216 to five (5) gallons of water.

GERMICIDAL INDICATOR: FORMULA 4216 is its own indicator of germicidal activity. In use dilution, it exhibits an amber color. Just as long as color remains in this solution, complete and rapid germicidal action is assured. Once color has disappeared, the solution should be discarded.

HOSPITAL USE: Use three (3) ounces of FORMULA 4216 to five (5) gallons of water (75 PPM titratable iodine) for thermometer disinfection, isolation and terminal disinfection procedures, disinfection of rubber sheets and rubber gloves, tubing and catheters. (5 minutes contact recommended.)

SURGICAL INSTRUMENTS: Remove all adhering blood and serous exudates by cleaning with a solution containing 3 ounces of FORMULA 4216 to 5 gallons of water. After cleaning, soak in a similar solution for at least 15 minutes.

SCHOOL USE: For the maintenance of public sanitation, it is suggested that FORMULA 4216 be employed for the cleansing and disinfection in one application of: walls, floors, desks, etc. as well as its utilization in shower rooms, locker rooms and washrooms.

GENERAL USE: This material should be used to clean (and disinfect) floors, walls, furniture, Venetian blinds, metal cabinets, drinking fountains, etc.

CAUTION: Harmful if swallowed. May cause skin and eye irritation. Avoid contact with skin, eyes or clothing. If in contact, wash thoroughly with water. If irritation persists, get medical attention. Avoid contamination of foods and feeds.

. Rinse empty container thoroughly with water and discard it.

7871