


WEST W fly bait

This product is authorized for use under USDA Poultry, Meat, Rabbit and Egg Products Inspection Programs.

ACCEPTED
Aug 16, 1971
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 52787 SUBJECT TO ATTACHED COMMENTS.



NET CONTENTS: 15 LBS

| | |
|--------------------------------------|---------|
| ACTIVE INGREDIENTS: | |
| 2,2-dichlorovinyl dimethyl phosphate | 0.93% |
| *Related Compounds | 0.07% |
| INERT INGREDIENTS: | |
| | 99.00% |
| | 100.00% |

DDVP
WEST FLY BAIT for control of resistant and non-resistant flies in Animal Buildings, Milk Sheds and Outdoor Areas. WEST FLY BAIT is a ready-to-use dry bait, or it may be dissolved in water at the ratio of thirteen (13) ounces of bait in one (1) gallon of water for use as a bait spray.

WARNING: Keep Out Of Reach Of Children. Do not take internally. Harmful if swallowed. See rear panel for additional warning.

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed.

DIRECTIONS FOR USE

DAIRY BARNs, MILK SHEDS, STABLES, LIVESTOCK BARNs, LOAFING SHEDS, PIG PENS, AND OUTDOOR AREAs—Scatter dry bait at rate of 1/4 pound per 1000 sq. ft., or apply bait spray at 1-3 gallons per 1000 sq. ft., as a coarse wet spray on floor or ground areas where flies congregate. Ideal baiting sites are around doorways, feed storage rooms, alleyways and window sills. On freshly limed floors, place treatment on feed sacks or paper. Repeat as necessary for control.

CAGED POULTRY AND TURKEYS ON WIRE—Apply 1/4 pound of dry bait or spray 1/2 gallon bait solution per 100 sq. ft. on droppings beneath cages or perches. Repeat weekly.

POULTRY HOUSES—Apply 1/4 pound dry bait or 1/2 gallon bait solution per 100 sq. ft. where flies congregate, but keep out of reach of birds. Apply to manure, window sills, outside of penned enclosures and on floors of feed storage room. Repeat as necessary.

DOG KENNELS—Apply 1/4 pound dry bait or 1-3 gallons of bait solution per 1000 sq. ft. outside dog runway, window sills, ledges and other surfaces out of reach of animals. For caged units apply to dropping area. Repeat as necessary.

To reduce fly populations in and around **OUTDOOR LIVING AREAs, PICNIC GROUNDS, BACKYARD AREAs, PATIOS, LOADING DOCKS, OUTDOOR LATRINES, PARKING AREAs, REFUSE AREAs, AROUND SERVICE STATIONS, OPEN-AIR DRIVE-INS, ICE CREAM STANDS, GARBAGE COLLECTION AND DISPOSAL AREAs**—Apply 1/2 pound dry bait or 1-3 gallons bait solution per 1000 sq. ft. of area. Apply on ground or on areas where flies congregate. For open air drive-ins and ice cream stands, pay particular attention to trash barrels and garbage cans where flies congregate.

WARNING: Keep Out Of Reach Of Children. Poultry and Domestic animals. Contains sugar, may be attractive to children. Do not take internally. Harmful if swallowed, inhaled or absorbed through the skin. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water, foodstuffs, milk or milking equipment. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.

2-PAM is also antidotal and may be administered in conjunction with atropine.

This product is toxic to wildlife. Birds feeding on treated areas may be killed. Apply only as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not reuse container. Destroy when empty.

4097-A 7715

Printed in U.S.A.

manufactured for
WEST
W

WEST CHEMICAL PRODUCTS, INC.
42-16 WEST STREET, LONG ISLAND CITY, N. Y. 11101
Branches in principal cities