

[52-193]

# X-TERM-O

## ACTIVE INGREDIENTS

- Petroleum Hydrocarbons..... 50.0%
- \*Technical Piperonyl Butoxide..... 20.0%
- \*\*N-Octyl Bicycloheptane Diethyl Carbamate..... 15.0%
- Pyrethrins..... 0.40%
- \*Equivalent to 0.40% of (1,1-dimethyl-3-methylbutyl) (6-Propylpiperonyl) Ether and 0.10% Related Compounds.

X-TERM-O is a concentrated insecticide for use with <sup>spray</sup> equipment. X-TERM-O is effective as a fog, contact spray, residual spray and has repellent action.

## FLYING INSECTS DIRECTIONS FOR USE

Exposed stages of FLIES, MOSQUITOES, GNATS, BEES, WASPS, HORNETS, INDIAN MEAL MOTHS, ANGIOUM'S GRAIN MOTHS, MEDITERRANEAN FLOUR MOTHS, FRUIT FLIES, CIGARETTE BEETLES, FLYING CLOTHES MOTHS.

To control these and similar small flying insects fog X-TERM-O at the rate of 1 ounce per 1000 cubic feet.  
Turn off all ventilating

equipment, close all windows and doors. Keep treated area closed for at least 10 minutes after treatment. For clothes moths repeat application monthly.

ACCEPTED  
JUL 14 1967  
52-193

## CRAWLING INSECTS DIRECTIONS FOR USE

Exposed stages of COCKROACHES, WATERBUGS, SILVERFISH, FIREBRATS, BOOKLICE, MILLIPEDES, CENTIPEDES, SOYBUGS, ANTS, TERMITES, SPIDER BEETLES, MEAL WORMS, CADELLES, CONFUSED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, SPIDERS, WEEVILS, DERMESIDS.

- For space spray — fog X-TERM-O at 2 ounces per 1000 cubic feet.
- As a contact spray — apply X-TERM-O directly on insects or wherever they are seen. Repeat as necessary.
- For flour and grain pests — clean all farinaceous material out of bins, elevator boots and spouts and apply X-TERM-O
- As a residual spray — apply X-TERM-O as a coarse wet spray at the rate of one gallon per 1000 square feet or to point of

run-off using a pressure sized sprayer. Thoroughly treat all crevices and breeding areas, cracks, crevices, under sides of work tables, behind sink and escutcheon plates, walls, window and door frames, elevator shafts and pits, behind and under equipment, laundry carts, garbage areas, warm dark moist areas, void spaces and shelves. Repeat as necessary.  
Repellent action may be developed by regular weekly spraying of area with X-TERM-O

**CAUTION: KEEP OUT OF REACH OF CHILDREN** Harmful if swallowed. Do not take internally. Avoid contamination of food. Food processing plants should be fogged when not in operation. Food should be removed or covered during treatment. Food processing equipment should be covered or thoroughly cleaned before using. Store away from heat or flames. Fabrics which are soiled or dusty may be stained by insecticidal fogs.

151-1-6-10      USDA Reg. No. 52-39      \*\*U.S. Patent No. 2,476,512      PRINTED IN U.S.A.

MANUFACTURED FOR

CHEMICAL TESTING CORP. 32-10 37th Avenue, Long Island City 1, N.Y.