

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER:

Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, promptly drink a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Induce vomiting. Call a physician immediately.

DIRECTIONS FOR USE:

GENERAL CLASSIFICATION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

BEFORE USING:

Contents of attached small container of activator must be added to a gallon Totacide-28. Shake well. Do not dilute. The final solution will exhibit a fluorescent yellow-green color.

Unused activated TOTACIDE-28 is effective up to 28 days. Used solutions should not be reused.

pH RANGE OF ACTIVATED PRODUCT: 6.9-7.4

For sterilization and disinfection of instruments such as lensed instruments, forceps, anesthesia equipment, and tubing, remove gross contamination and clean thoroughly. Rinse and damp dry objects before immersing in undiluted TOTACIDE-28 to which activator has been added. Use covered containers to minimize evaporation or odor and prevent contamination. Follow authorized aseptic hospital procedures for flushing instruments before use.

4-A0181

TO STERILIZE:

Completely immerse previously cleaned objects in undiluted TOTACIDE-28, to which activator has been added, for a minimum of ten (10) hours to destroy resistant pathogenic spores of *Clostridium sporogenes*, *Bacillus subtilis*, *Clostridium tetani* and *Clostridium perfringens*.

TO DISINFECT:

Completely immerse previously cleaned objects in undiluted TOTACIDE-28 to which activator has been added, for a minimum of ten (10) hours to destroy *M. tuberculosis*, *Pseudomonas aeruginosa*, *Trichomonas vaginalis*, *Leptospira* and the *Virus of Herpes simplex*, *Varicella*, *Adenovirus*.

NOTE:

When removing objects from TOTACIDE-28 solution use aseptic technique and RINSE OBJECTS THOROUGHLY, inside and out, with sterile water prior to use.

Do not immerse carbon steel objects over 24 hours. TOTACIDE-28 solution contains a rust inhibitor but will not prevent or remove rust caused by plating or improper care of instruments.

STORE IN COOL PLACE

Mixing Date _____ Expiration Date _____

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Unused material or rinsate should be disposed of in landfill approved for pesticides or buried in safe place away from water supplies.

Triple rinse container and dispose of in incinerator or approved landfill or bury in safe place. Consult Federal, state or local authorities for approved alternate procedures.

MADE IN U.S.A.

NET
CONTENTS:
1 GALLON
U.S.
(3.785
LITERS)

WE
MANUFACTURED
WEST



FOR INSTITUTIONAL AND PROFESSIONAL USE ONLY

tōtacide-28TM

SPORICIDAL, TUBERCULOCIDAL, BACTERICIDAL, VIRUCIDAL*,
FUNGICIDAL, COLD STERILIZING AGENT. RECOMMENDED FOR USE
IN HOSPITALS FOR THE STERILIZATION AND DISINFECTION OF:

• INSTRUMENTS • ANESTHESIA EQUIPMENT • LENSED INSTRUMENTS • THER-
MOMETERS • RUBBER OBJECTS • PLASTIC OBJECTS

ACTIVE INGREDIENT:

Glutaraldehyde 2.0%

INERT INGREDIENTS:

..... 98.0%

U.S. Patent No. 4,093,744

Keep Out Of Reach Of Children.

IRRITATING TO EYES

DANGER:

(See back panel for First Aid Statement and
additional precautionary statements.)

NET
CONTENTS:
1 GALLON
U.S.
(3.785
LITERS)



MANUFACTURED BY

WEST CHEMICAL PRODUCTS, INC., 1200 Hill of Town Road, Princeton, N.J. 08540

EPA Reg. 52-252

ACCEPTED

MAR 16 1981

Under the provisions of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 52-252

138

DIRECTIONS: Dissolve contents of this vial into one gallon of TOTACIDE-28. Activated solution will be fluorescent yellow-green.

STORE IN COOL PLACE


DANGER: Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, promptly drink a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Induce vomiting. Call a physician immediately.

440181

MADE IN U.S.A.



Aqueous Totacide-28 Activator

for use with Totacide-28.

INGREDIENTS: Contains Tripotassium Phosphate, D & C Green #8; Potassium Nitrite and water.

U.S. PATENT NO. 4,093,744

DANGER:
Keep Out Of Reach Of Children.
Harmful if swallowed.
(See back panel for additional dangers.)

NET CONTENTS: 30 gms.



MANUFACTURED FOR:
WEST CHEMICAL PRODUCTS, INC.
1000 Herrontown Road,
Princeton, New Jersey 08540