



000056-00023-121184

Printed: 12:49:58 Friday, 21 May, 1999 # 16 / 2882

Systems Integration Group, Inc.

[FRONT PANEL]

EATON'S <sup>®</sup>  
"SEMI-PERMANENT"  
BAIT BLOCKS <sup>®</sup>  
RODENTICIDE

ACCEPTED  
DEC 11 1984  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 56-23

ACTIVE INGREDIENT:  
Diphacinone (2 diphenylacetyl-1,  
3-indandione) 0.005%  
INERT INGREDIENTS: 99.995%  
100.000%

CAUTION  
KEEP OUT OF REACH OF CHILDREN  
SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

This Package Is Not to Be Broken for Resale of Enclosed Units

MFG. BY

J. T. EATON & CO., INC.  
1393 East Highland Road  
Twinsburg, Ohio 44087

EATON'S <sup>®</sup> and BAIT BLOCKS <sup>®</sup> are Registered Trademarks of J. T. EATON & CO., INC.

EPA REG. No. 56-23  
EPA EST. No. 56-OH-1

Contents: 40 - 8 ounce BLOCKS

NET WT. - 20 POUNDS

[LEFT PANEL]

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND  
DOMESTIC ANIMALS

CAUTION - Keep away from humans, domestic animals and pets.  
If swallowed, this material may reduce the clotting ability  
of the blood and cause bleeding.

NOTE TO PHYSICIAN - If ingested, administer Vitamin K<sub>1</sub>  
intramuscularly or orally, as indicated in bishydroxycoumarin  
overdoses. Repeat as necessary based on monitoring of prothrombin  
times.

ENVIRONMENTAL HAZARD

Keep Out of Lakes,  
Streams or Ponds.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product  
may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty box into application equipment.  
Then dispose of box in a sanitary landfill or by incineration, or, if  
allowed by state and local authorities, by burning. If burned, stay  
out of smoke.

NOTICE: Buyer assumes all risks of use, storage or  
handling of this material not in strict accordance  
with directions given herewith.

[RIGHT PANEL]

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS - Eaton's "Semi-Permanent" BAIT BLOCKS® can be used in areas where they will not spoil because of excessive amounts of moisture or mildew (as with a cereal-type bait) such as:

1. In food processing plants (dairies, meat packing plants, sausage works, etc.) where the work rooms are washed down each night.
2. Outdoors in animal feed lots, along side of grain elevators, grain storage bins, barns, etc.
3. Attached to pier pilings, wharf runways, along breakwalls, etc.
4. Along gullies, irrigation ditches, rivers, sewers, etc.
5. Along railroad sidings, clean-out yards, etc.
6. In and around dumps, landfills, etc.
7. Under loading docks, along fences, outdoor storage areas, etc.

Treated baits must be placed in locations not accessible to children, pets, domestic animals, wildlife or in tamper-proof bait boxes. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS - Determine areas where rats will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

APPLICATION DIRECTIONS

INDOOR USE - Secure a BAIT BLOCK® to one end of a tie-wire and attach other end to a pipe, piece of machinery, etc...in areas where rats are known to feed, water, run or exist - EATON'S BAIT BLOCKS® will do the rest

OUTDOOR USE - Secure a BAIT BLOCK® to one end of a tie-wire and attach other end to a fixed object (wood post, pipe, etc.) in an area where rats are known to feed, water, run or exist - EATON'S BAIT BLOCKS® will do the rest.

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.