



# EATON'S A-C FORMULA 50

Ready-To-Use **RODENTICIDE**

No Prebaiting Required - No Tolerance Build-up - Just Use as Directed.

**SUGGESTED USE:** Place EATON'S A-C FORMULA 50 in protected areas where Rats and Mice are known to feed, water, run or exist.

- ... Use shallow containers for the bait and protect bait from children, pets and domestic animals.
- ... For RATS, use one or two bait placements (¼ to 1 pound each) at every point of activity.
- ... For MICE, tablespoon amounts (¼ to ½ ounce each) of bait should be placed at 8-12 foot intervals in the infested area. Baits should be placed in locations not accessible to children, pets and domestic animals, or in tamperproof bait boxes.

**KEEP FEEDING STATIONS WELL SUPPLIED:** Inspect and replenish bait stations daily for the first week and regularly thereafter until the bait is no longer taken, which usually takes from 5-14 days. Never allow the feeding

station to remain empty for over 48 hours. Sour, moldy or otherwise decomposed bait should be replaced promptly.

**CAUTION:** When fresh meat or fresh fish is added to this bait, precaution should be taken to prevent feeding by dogs or cats. These animals have been known to actively search out baits of this type. Therefore, baits must be placed so as to prevent feeding by these animals. If eaten by humans, cats and domestic animals it may reduce the clotting ability of the blood and cause bleeding. In such case, blood transfusion and Vitamin K are indicated, as in a case of hemorrhage caused by an overdose of bishydroxycoumarin.

**ACTIVE INGREDIENT:**

2-Pivalyl-1, 3-Indandione

0.025%

**INERT INGREDIENTS:**

99.975%

100.000%

\_\_\_\_\_ LBS. NET WEIGHT

FOR PROFESSIONAL & PUBLIC HEALTH USE ONLY

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

USDA Reg. No. 56-27

Manufactured and Controlled by