PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN — If ingested, administer Vitamin K1 intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD

Keep Out of Lakes, Streams or Ponds.

DIRECTIONS FOR USE


It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DISPOSAL

DO NOT REUSE EMPTY CONTAINER.
WRAP CONTAINER AND PUT IN TRASH
FOR COLLECTION

SELECTION OF TREATMENT AREAS

Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.



ACTIVE INGREDIENTS:
Diphacinone [2-(diphenylacetyl)-1, 3-indanedione]
Diphacinone, Sodium Salt [2-(diphenylacetyl)-1,
3-indanedione, sodium salt]
INERT INGREDIENTS:

0.0025%

0.0027% 99.9948% 100.0000%

CAUTION:

KEEP OUT OF REACH OF CHILDREN SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

J.T. EATON & CO., INC.

1393 East Highland Road • Twinsburg, Ohio 44087 EATON'S® 3 BAI's BLOCKS® are Registered Trademarks of J.T. EATON & CO., INC. FLAVORIZER'™ is a Trademark of J.T. EATON & CO., INC.

NET WT. 10 LBS.

EPA EST. NO. 56-OH-1 EPA REG. NO. 56-42

SPECIAL FORMULA

EATON'S* All-Weather 3AIT BLOCKS*
Rodenticide contains selected inert ingredients and 2 anti-coagulants blended into a ready-to-use palatable rodenticide that is lethal to rats when consumed over a period of several days.

SUGGESTED USE

For use in wet or damp areas, snap off EATON'S® All-Weather BAIT BLOCKS® Rodenticide into the handy marked squares and place into protected areas where rats or mice are known to feed, water or exist.

APPLICATION DIRECTIONS

NORWAY RATS AND ROOF RATS: Apply 2 to 8 squares (4 to 16 oz.) of bait per placement. Place bait at 15 to 30 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Break each 2 oz. square into four or more pieces. Piece bait at 8 to 12-foot intervals. Use one-fourth to one-half of a broken square at each piacement site for light to moderate infestations. Where mouse activity is extremely high, the material from one broken square should be placed at each placement site. Larger placement (up to 2 oz.) may be be needed at points of very high mouse activity. Maintain an unterrupted supply of fresh bait for at least 15 days.

RATS & MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replanish bait as needed.