


PM 16 56-56 1A3

FRONT



**EATON'S A-C FORMULA 90  
READY-TO-USE RODENTICIDE**

**No Prebaiting Required**

**Just Use As Directed**

**ACTIVE INGREDIENT:**

Chlorophacinone  
2-[(p-chlorophenyl) phenylacetyl],  
3-indandione 0.005%

**INERT INGREDIENTS:** 99.995%  
100.000%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS**

**Rodents may consume a lethal dose in one feeding with first  
dead rodents appearing four or five days after treatment  
begins.**

**MFG. BY**

**J. T. EATON & CO., INC.  
1393 EAST HIGHLAND ROAD  
TWINSBURG, OHIO 44087**

**EPA REGISTRATION NO. 56-56  
EPA EST. NO. 56-OH-1**

**NET WT.**

2.8 f 3

**BACK PANEL**

**DIRECTIONS FOR USE**

**It is a violation of federal law to use this product in a manner inconsistent with its labeling.**

**USE RESTRICTIONS:** For control of Norway Rats, Roof Rats and House Mice in and around homes, industrial and agriculture buildings, and similar man-made structures.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

**SELECTION OF TREATMENT AREAS -** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain and snow. Remove as much food as possible.

**APPLICATION DIRECTIONS:**

**RATS:** Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

**MICE:** Apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

**RATS & MICE:** Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and uneaten bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

**BEST AVAILABLE COPY**

RIGHT PANEL

PRECAUTIONARY STATEMENTS

CAUTION

HAZARD TO HUMANS AND  
DOMESTIC ANIMALS

CAUTION - Keep away from humans, domestic animals and pets. If swallowed, this material may reduce clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K1 intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD

Keep Out of Lakes,  
Streams or Ponds.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Dispose empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**BEST AVAILABLE COPY**