

# Saponified Cresylic Solution

## For Disinfecting Stables, Sick Stalls

Phenol Coefficient 3--A.O.A.C. Method

### Active Ingredients

Cresylic Acid.....	50%
Soap.....	23%
Inert Ingredients.....	27%
	100%

**DANGER**  **POISON:** Keep out of reach of children.

See antidote statement and precautions on back panel. Read entire label carefully before using.

**Net Contents One Gallon (U.S. Measure)**

EPA Reg. No. 59-7

## Saponified Cresylic

### DIRECTIONS FOR USE

#### Dilution Rate

Use 1 part of Saponified Cresylic Solution to 10 parts of water.

#### Disposal

Rinse empty container thoroughly and discard.

#### GENERAL DISINFECTION (Livestock)

Saponified Cresylic Solution is used in disinfecting stables, barns, cars, boats, and other vehicles as transport vehicle, affected with infectious, or communicable disease.

Remove all livestock from building treated with Saponified Cresylic.

Thoroughly clean and protect the building, all equipment, and water. Remove all litter from the building thoroughly.

Apply Saponified Cresylic Solution to floors, walls, and equipment and thoroughly disinfecting equipment as to remove scrub with water and brush.

Ventilate the building, and after antiseptic treatment has been applied.

Do not use in sick animals, and make sure they are not treated.

Keep all pipes, floors, and equipment washed with Saponified Cresylic to reduce the spread of disease.

#### Tuberculosis

Saponified Cresylic Solution is a permitted fungicide for use in the control of Tuberculosis, as authorized by the Department of Agriculture, U.S. Department of Agriculture.

# For Disinfecting Stables, Sick Stalls

Phenol Coefficient 3—A.O.A.C. Method

<b>Active Ingredients</b>	
Cresylic Acid.....	50%
Soap.....	23%
Inert Ingredients.....	27%
	<hr/>
	100%



**DANGER POISON:** Keep out of reach of children.

See antidote statement and precautions on back panel. Read entire label carefully before using.

**Net Contents One Gallon (U.S. Measure) EPA Reg. No. 59-7**



infectious or communicable disease.  
 Remove all manure from stall treated with Saponified Cresylic Solution.  
 Thoroughly clean and scrub the building and equipment with water. Remove all dirt from the building floor.  
 Use disinfectant solution to apply it to floors, walls, ceiling equipment until thoroughly wet. Watering equipment is. Thoroughly scrub with detergent and rinse.  
 Ventilate the building and air until treatment has been absorbed.  
 Do not use in milking stalls, in milk houses or poultry houses.  
 Keep your premises and equipment disinfected with Saponified Cresylic to reduce the hazard of disease infection.

### Tuberculosis

Saponified Cresylic Solution is permitted for use by the Army and Veterinary Service, Animal & PHS Service, U.S. Department of Agriculture disinfection program, and all other housing and treatment facilities directed under General Disinfectant

CUST. <u>W. Cooper</u>		DESIGN <u>SAPONIFIED CRESYLIC SOLUTION</u>		NO. PROOFS <u>10</u>		ARTIST <u>B. C.</u>
ENG. <u>2276-2</u> S.O. <u>242-7693</u>		SIZE <u>402x610x907</u> C.E.A. <u>562x20.400</u> B.P. <u>64-201</u> DATE <u>11-14-72</u>		PROOF DATE <u>11-29-72</u>		
CHANGE A <u>3-6-73</u>		CHANGE B		CHANGE C		
CHANGE AS DETAILED <input type="checkbox"/>		APPROVED AS SUBMITTED <input type="checkbox"/>		CUSTOMER SIGNATURE		
				DATE		

READ BEFORE SIGNING: Customer agrees that (1) Continental Can Company, Inc. shall not be liable for (a) any error in the attached unless this proof is returned to Continental with such error plainly noted in writing, or (b) the failure of any part of the attached to comply with any local, state or Federal law or regulation, or (c) any part of the attached which infringes the property rights of others acquired without limitation thereof, by assignment, trademark, copyright, letters patent or otherwise and (d) production colors not being exactly the same as the hand proved colors, if any, in the attached, although Continental will duplicate the colors as closely as possible. (2) Customer shall indemnify and hold Continental harmless against the liability described in (1) above, including attorney fees and court costs or for consequential damages, and (3) the liability of Continental for uncorrected errors plainly noted in the attached shall not exceed the actual cost of the containers bearing the noted error. Changes in these costs and delay priority from CONTINENTAL CAN COMPANY, INC.

# Saponified Cresylic Solution

## DIRECTIONS FOR USE

### Dilution Rate

Use 4½ fluid ounces of Saponified Cresylic Solution to one gallon of water.

### Disposal

Rinse empty container thoroughly with water and discard.

## GENERAL DISINFECTION (Livestock)

Saponified Cresylic Solution is recommended for use in disinfecting stables, barns, kennels, cattle cars, boats and other vehicles used to house or transport livestock affected with any contagious, infectious, or communicable disease.

Remove all livestock from buildings to be treated with Saponified Cresylic Solution.

Thoroughly clean and polish or remove from the building all equipment, including feeders and waterers. Remove all litter or bedding and clean the building thoroughly.

Use disinfectant only at recommended rate, and apply it to floors, walls, ceilings, and livestock equipment until thoroughly wet. After feeding and watering equipment is disinfected, thoroughly scrub with detergent and rinse with potable water.

Ventilate the building and do not house livestock until treatment has been completed, wet or dried.

Do not use in milking stables, milking parlors, milk houses or poultry houses.

Keep your premises and equipment clean, and disinfect with Saponified Cresylic Solution to reduce the hazard of cross infection.

## Tuberculosis

Saponified Cresylic Solution is a tuberculicide permitted for use by the Animal Health Program, Veterinary Service, Animal & Plant Health Service, U.S. Department of Agriculture, in official

## Brucellosis (Contagious Abortion)

Saponified Cresylic Solution is a disinfectant recommended for cleaning and disinfection of premises after Brucellosis infections have been removed from the herd. Use as suggested under General Disinfection.

## Hog Cholera

Pens and equipment should be disinfected as directed under General Disinfection.

## Official Meat Processing Establishments

Saponified Cresylic Solution has been approved by the Meat and Poultry Inspection Program, Animal and Plant Health Inspection Service, U.S. Department of Agriculture, for use in tarrets and/or dressing rooms of meat processing plants.

## INFECTIOUS DISEASES (Human)

Saponified Cresylic Solution should be used to disinfect everything which comes into contact with the patient (including bedding, utensils) at the recommended rate. Before treated food contact surfaces with potable water before reuse.

## Important

Always remove all litter and feces before applying disinfectants.

## DANGER

May be fatal if swallowed or absorbed through the skin. May produce severe burns if contact with eyes, or irritation of clothing. Avoid breathing spray mist. Wash thoroughly after handling. Use this product only as directed on the label.

## POISON—ANTIDOTE

Human Intoxication: Wash immediately with copious amounts of water. If swallowed, induce vomiting and give activated charcoal or milk if available. For eye contact, flush with water for at least 15

transport livestock affected with any contagious, infectious, or communicable disease.

Remove all livestock from buildings to be treated with Saponified Cresylic Solution.

Thoroughly clean and protect livestock from the building, all equipment, including feeders and waterers. Remove all litter or bedding and clean the building thoroughly.

Use disinfectant only at recommended rate, and apply it to floors, walls, ceilings, and livestock equipment until thoroughly wet. After feeding and watering equipment is disinfected, thoroughly scrub with detergent and rinse with potable water.

Ventilate the building and do not house livestock until treatment has been absorbed, set or dried.

Do not use in milking stalls, milking parlors, milk houses or poultry houses.

Keep your premises and equipment clean, and disinfect with Saponified Cresylic Solution to reduce the hazard of cross infection.

## Tuberculosis

Saponified Cresylic Solution is a tuberculocide permitted for use by the Animal Health Programs, Veterinary Service, Animal & Plant Health Service, U.S. Department of Agriculture, in official disinfection programs involving domestic animal housing and transportation facilities. Disinfect as directed under General Disinfection.

Department of Agriculture, for use in stables and for dressing rooms of meat processing plants.

## INFECTIOUS DISEASES (Human)

Saponified Cresylic Solution is used to prevent and control everything from the contact with the patient to the final disposal of the body and bedding. For a complete list of contact surfaces, turn page 2 over before use.

### Important

Always remove all litter and filth before applying disinfectant.

## DANGER

May be fatal if swallowed or absorbed through the skin. May produce severe burns. Do not get in eyes, on skin or on clothing. Avoid breathing spray mist. Wash thoroughly after handling. Use this product only as directed on the label.

## POISON—ANTIDOTE

Human-Internal: If swallowed, swallow immediately, and administer 2 or 3 table-spoons of olive oil, induce vomiting and give 1 quart water. alburnous drink, or milk or water. If any vomit. External: In case of contact, immediately wash eye with running water for 15 minutes. For eye, get to hospital as quickly as possible. Remove contact lenses if they are present. Wash and launder before reuse.

## COOPER U.S.A., INC.

Cooper Building, P.O. Box 1000

A Wellcome Company

10	ARTIST
10-6-73	