

Cooper-Tox Extra

DIRECTIONS FOR USE

Beef Cattle

Ticks, Horn Flies and Lice—Accurately mix Cooper-Tox Extra at the rate of 1 gallon to 150 gallons of water for complete kill of ticks, horn flies and lice present, and for protection against reinfestation. Thorough wetting of animals is essential for best results. Repeat application as necessary.

Back Rubbers—To aid in controlling horn flies and lice on beef cattle, dilute 1 pint Cooper-Tox Extra with 1½ gallons diesel fuel or light motor oil. Charge burlap roll by soaking with solution so it is thoroughly wet, but not to the point of dripping. Repeat charging as necessary. This application results in rapid reduction of horn flies after exposure to animals, but regular exposure for about ten weeks is usually necessary for good results against lice. Do not treat animals under six months of age or within 30 days of slaughter.

Sheep and Goats

Lice and Sheep Ticks (Keds)—Mix Cooper-Tox Extra at the rate of 1 gallon to 500 gallons of water. Thorough wetting is essential for best results. A single application is usually sufficient for control of these pests.

Sheep Scab (Psoroptic Mange)—Accurately mix Cooper-Tox Extra at the rate of 1 gallon to 100 gallons of water for complete cure of scab in a single dipping. Immerse sheep not less than ½ minute, ducking heads at least twice. When large bands are dipped, the dip should be frequently checked by a vatside test to assure proper dipping strength.

Fleece Worms and Ticks—Dilute Cooper-Tox Extra at the rate of 1 gallon to 150 gallons of water. Wet sheep thoroughly. Repeat as necessary.

Hogs

Lice—Mix Cooper-Tox Extra at the rate of 1 gallon to 150 gallons of water. A single application is usually sufficient to control this pest. Do not treat sows within two weeks of farrowing nor at least three weeks thereafter.

Sarcoptic Mange—Mix Cooper-Tox Extra at the rate of 1 gallon to 100 gallons of water. Repeat application in 14 days. Thorough wetting of animals is essential. Do not treat sows within two weeks of farrowing nor for three weeks thereafter.

For Smaller Quantities

2 tablespoons of Cooper-Tox Extra makes 1 gallon of spray or dip.

Residual Surface Spraying

Flies, Mosquitoes—Mix Cooper-Tox Extra at the rate of 1 gallon to 40 gallons of water. Wet surfaces thoroughly, but not to the point of excessive run-off. Spray outside walls of dairy barns, on fences, inside hog pens, on garbage cans and similar places. Repeat as needed.

Mixing Cooper-Tox Extra with Water

Add Cooper-Tox Extra directly to required amount of water in accordance with dilutions recommended on this label. Stir or agitate the resulting mixture thoroughly before each use.

Replenishment (Dipping Vats)

Add Cooper-Tox Extra in accordance with the recommended dilutions for all fresh water added to the vat, or control replenishment by vatside test.

WARNING

May be fatal if swallowed. Cooper-Tox Extra is toxic and is absorbed through the skin in dry form and from solutions. Do not get in eyes, on skin or on clothing. Do not breathe spray mist. Use respirator or goggles for protection if necessary. Keep away from food and food products. Keep clothing free from residue. Wash hands, arms and face with soap and water after handling and before eating or smoking. Concentrations of Cooper-Tox Extra greater than those recommended should never be used.

Do not use on dogs, cats, or any other household pets.

Do not use on any animals under 3 months of age. Do not dip animals under 6 months of age. If treatment of young animals is necessary, spraying is recommended. Avoid spraying or dipping animals in cold, stormy weather, and avoid treating overheated or sick animals. Do not use Cooper-Tox Extra on dairy animals, in dairy barns or milk rooms, in poultry houses, or on food to be fed to livestock. Do not spray within 30 days or dip within 60 days of slaughter.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

Do not use, pour, spill or store near heat or open flame.

Antidote

Human-Internal: Call a physician immediately. Give sodium chloride 15 grams (one tablespoonful of salt) in a glass of warm water. Repeat until vomit is clear. **External:** Wash with plenty of soap and water.

COOPER U.S.A., INC.

Research Triangle Park, N.C. 27709 USA
A Wellcome Company

Q21870

Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 5-9 5 5