

{59-67}

50% Malathion

Emulsifiable Concentrate

For Home-Farm-Ranch

Active Ingredients

Malathion*

Petroleum Distillates

Inert Ingredients

50.0

45.0

5.0


CAUTION: Keep out of reach of children. See other cautions on back panel.

Net Contents One Gallon (U.S. Measure)

EPA Reg. No. 59-67



COOPER



50% Malathion Emulsifiable Concentrate

DIRECTIONS FOR USE

Poultry- For Lice and Northern Fowl Mite

Dilute 50% Malathion at the rate of 2 teaspoons in 1 gallon of water (1 to 100). Spray 1 gallon of mixture over 1 sq. ft. of area. Repeat in 4-8 weeks if necessary.

Poultry House- Northern Fowl Mite, Chicken Red Mite, Lice and Flies

Dilute 50% Malathion at the rate of 4 tablespoons in 1 gallon of water (1 to 50). Apply thoroughly to walls, ceiling, cracks, other nooks and crevices. Make sure to force spray into cracks and crevices. Repeat if necessary.

Dogs & Cats- Fleas

Dilute 1 ounce of concentrate in 1 gallon of water. Spray on damp, wet pet thoroughly. Repeat every 2-3 days if necessary. Do not treat animals under 1 month of age.

Fleas in Kennels, Pens, Yards, Lawn and Under Houses

Use a spray containing 1 ounce of 50% concentrate in 1 gallon of water. Apply to kennels, pens, yards, lawns, etc. Repeat if necessary. Do not spray around or on plants or animals. Do not spray around or on children. Do not spray on or near food.

House Flies- Dairy Barns, Hog Barns, Poultry Houses, Calf Pens, Manure Piles, Garbage Cans, Yards and around Meat Processing Plants

Use a spray containing 1 ounce of 50% concentrate in 1 gallon of water. Apply to manure piles, garbage cans, yards, etc. Repeat if necessary. Do not spray around or on plants or animals. Do not spray around or on children. Do not spray on or near food.

Vegetables

Dilute 1 ounce of concentrate in 1 gallon of water. Spray thoroughly over the vegetables. Do not spray on or near food. Do not spray on or near children. Do not spray on or near pets. Do not spray on or near plants or animals. Do not spray on or near food.

Ornamentals

Dilute 2 teaspoons of 50% Malathion in 3 quarts of water for control of aphids, scale insects, leaf miners, bag worms and spider mites on firs, yews, spruce, arborvitae, boxwood and other ornamentals. Injury may occur to certain ferns including Maidenhair Ferns and Boston as well as some species of Camellia.

Flowers

Dilute 2 teaspoons of 50% Malathion in 3 quarts of water for control of aphids, plant bug, leaf hoppers, scale insects and white flies on gladioli, roses, phlox and gladiolas.

Fruit Trees

Dilute 1 tablespoon in 3 quarts of water for control of scale insects, bud moth, spider mites, woolly aphids, mealy bugs and leaf rollers on apple, cherry, pear, plum and apricots. Do not apply within 3 days of harvest and 7 days before harvest of apricots.

CAUTION

Harmful if inhaled, swallowed or absorbed through the skin. Avoid breathing it. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after using. Do not use near food or drink.

Do not use on lawns or around milk or milk products. Avoid contact with children and pets. Do not use on or near food. Do not use on or near children. Do not use on or near pets.

This product is toxic to fish. Keep it out of wells, streams, ponds, etc. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on the label.

Do not use empty container. Do not use product after expiration date and do not use if label is missing.

COOPER U.S.A., INC.

Chicago, Illinois 60614 USA

A Wellcome Company