

Sugar Bait

Fly Killer

For Fast Fly Control
New With Stabilized DDVP
Use Wet or Dry

Active Ingredients


2,2-Dichlorovinyl dimethyl phosphate 0.465%
 Related Compounds 0.035%

Inert Ingredients 99.500%
 100.000%

CAUTION: Keep out of reach of children.
 See other cautions on back panel.

Net Contents One Pound

USDA Reg. No. 59-75



Sugar Bait Fly Killer

DIRECTIONS FOR USE

Recommended for dairy barns, poultry houses, livestock barns and pens. Kills both resistant and non-resistant flies. Can be used as a liquid or dry bait. Either use will give fast reduction of fly populations.

Dry Bait

Scatter Cooper Sugar Fly Bait at the rate of ¼ lb. per 1000 sq. ft. in and around areas where flies are seen. Apply every 2-3 days until fly population is under control and then apply as needed. Place baits in areas inaccessible to children and pets.

Liquid Bait

Dissolve 1 lb. of Sugar Fly Bait in 1 gal. of water. Sprinkle or spray floor areas, window sills, walls or other areas where flies are seen. Repeat daily until flies are under control and then apply as needed.

CAUTION

Harmful if swallowed. Induce vomiting if swallowed. Avoid prolonged breathing of dust and repeated or prolonged contact with skin. Wash hands after use. Keep out of reach of children, poultry and domestic animals. Avoid contamination of feed or foodstuffs. Do not use inside homes. Do not contaminate milk or milking equipment. Application in dairy barns should be made after milking.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Apply this product only as specified on this label.

COOPER U.S.A., INC.
 Chicago Illinois 60614 USA
A Wellcome Company