

Specifically Prepared for Use on Dogs, Cattle, Horses, Sheep, Goats and Swine as a Spray or Dip.

Kills Ticks, Fleas, Lice, Horn Flies, Keds and Wool Maggots.

Also Helpful in Controlling Screwworm Infestations.

No Waiting Period Between Last Application and Slaughter of Livestock.

Active Ingredients


Dioxathion [2,3-p-dioxanedithiol S, S-bis (O,O-diethyl phosphorodithioate)]	20.40%
Related Compounds	8.70%
Aromatic Petroleum Distillates	56.62%
Inert Ingredients	14.28%
	<hr/>
	100.00%

WARNING: Keep out of reach of children. See other warnings on back panel.

Net Contents One Gallon (U.S. Measure)

EPA Reg. No. 59-113-AA

SOLD TO VETERINARIANS ONLY



Co-Nav

DIRECTIONS FOR

DOGS

TICKS, FLEAS AND LICE
1 gallon of water. Spray

Repeat only when needed.

PREMISE SPRAY FOR

with 1 gallon of water.
Use to spray possible
and under various pieces

CATTLE, SHEEP, GOATS

TO KILL TICKS, HORNS

Mix at the rate of 1 gallon
required but no more
than a bucket of water

LONG-LASTING PROTECTION

For up to 3-week
protection and kill of
fleas and ticks. Wash up
and out of the hair.

SPRAYING: 1 gallon of

water per animal

Co-Nav

DIRECTIONS FOR USE

DOGS

TICKS, FLEAS AND LICE: Thoroughly mix at the rate of ½ fl. oz. (1 tablespoonful) with 1 gallon of water. Sponge, swab, bathe or dip dog making sure dog is thoroughly wet.

Repeat only when necessary but not more frequently than once every week.

PREMISE SPRAY FOR YARDS, KENNELS: Mix at the rate of 2½ fl. ozs. (5 tablespoonfuls) with 1 gallon of water. Use 1 gallon of spray per 1000 sq. ft. of lawn or kennel area. Make sure to spray possible hiding places such as cracks in cement or wood, around dog's bedding and under various pieces of debris.

CATTLE, SHEEP, GOATS, SWINE, AND HORSES

TO KILL TICKS, HORN FLIES, LICE, KEDS, WOOL MAGGOTS AND SCREWORMS: Mix at the rate of 1 gallon Co-Nav in 200 gallons of water. Mix thoroughly. Spray or dip as required but no more often than every two weeks. The use of Co-Nav requires no waiting period between last application and slaughter.

LONG-LASTING PROTECTION: Co-Nav provides long-lasting protection against reinfestation of ticks and 3-week protection against horn flies. A single application is usually sufficient for lice and keds. For biting lice on goats, a second application after 21 days may be required for complete clean-up. Co-Nav also aids in controlling screwworm larvae and provides up to 2 months protection against wool maggots.

SPRAYING: Make sure spray is thoroughly mixed before application. Wet animals thoroughly for best results. Do not allow animals to drink from run-off pools.

DIPPING

Drunk, ill or
malnourished
animals should
not be dipped.

REPLENISHMENT

Replenish dipping water with fresh water every 24 hours.

WARNING

Co-Nav is toxic and may be fatal if swallowed or absorbed through the skin. Do not inhale spray, get in eyes, on skin or on clothing. Use respirator and goggles if necessary. Wash with soap and water in case of contact.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

Do not use on dairy animals or in dairy barns. Do not dip or spray foals. Do not dip animals under 3 months of age.

Do not contaminate feed or foodstuffs. Food utensils such as teaspoons and tablespoons should not be used for food purposes after use with pesticides. Do not reuse container. Destroy when empty.

This product is toxic to fish, wildlife and birds. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

COOPER U.S.A., INC., Veterinary Products Division
Research Triangle Park, N.C. 27709

A Wellcome Company

V023080