

Co-Nav

RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS
OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES
COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

Specifically Prepared for Use on Dogs, Cattle, Horses, Sheep, Goats and Swine as a Spray or Dip.
Kills Ticks, Fleas, Lice, Horn Flies, Keds and Wool Maggots.
Also Helpful in Controlling Screwworm Infestations.
No Waiting Period Between Last Application and Slaughter of Livestock.

Active Ingredients	
Dioxathion (2,3-p-dioxanedithiol S,S-bis-(O,O-methyl phosphorodithioate))	20.40%
Related Compounds	7.10%
Heavy Aromatic Naphtha	53.50%
Inert Ingredients	19.00%
	100.00%

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.



Net Contents (One Gallon (U.S. Measure))

EPA Est. 59-103-1
EPA Reg. No. 59-113



DE
PLICATORS
FOR THOSE WHO
CARE.

and Same as a Spray or Dip.

20.40%
7.10%
53.50%
19.00%
100.00%

EN
PLACEMENTS
EPA Est. 59-NC-1
EPA Reg. No. 50.123

AC
Muda
Fung
in em
pestic
EPA R

Co-Nav

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS:
WARNING

... ..

STATEMENT OF PRACTICAL TREATMENT

... ..

ENVIRONMENTAL HAZARDS

... ..

PHYSICAL OR CHEMICAL HAZARDS

... ..

DIRECTIONS FOR USE

... ..

PREMISE S
Use 1/4-1/2
... ..

TO KILL TH
... 200 gal
of Co-Nav
LONG LAST
... ..
SPRAYING:
... ..
DIPPING:
... ..
REPLENISH
ment by 20

PROMI
Do not
PESTIC
... ..
CONTA
... ..
GENER

WELCOM
Burrough
Research
... ..

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (& DOMESTIC ANIMALS)
WARNING
Irritant if swallowed or absorbed through the skin. Do not inhale spray, get in eyes, use respirator or goggles if necessary. Wash with soap and water in case of contact. Do not use in dairy barns. Do not dip or spray foals. Do not dip animals under 3 months of age.
Do not use on foodstuffs. Food utensils, such as teaspoons and tablespoons should not be used with pesticide.
STATEMENT OF PRACTICAL TREATMENT
If swallowed, contact a veterinarian immediately. DO NOT INDUCE VOMITING unless under medical supervision. Cause aspiration pneumonia.
If in eyes, flush with plenty of soap and water.
If on skin, wash with plenty of soap and water.
If on clothing, wash with plenty of soap and water.
ENVIRONMENTAL HAZARDS
Toxic to fish and wildlife and birds. Keep out of any body of water. Do not contaminate water by disposal of wastes. Do not apply where runoff is likely to occur.
PHYSICAL OR CHEMICAL HAZARDS
None known.

DIRECTIONS FOR USE
Use this product in a manner consistent with its labeling.
Thoroughly mix at the rate of 1 fl. oz. (1 tablespoonful) with 1 gallon of water. Make sure dog is thoroughly wet.
Repeat treatments as directed on label.
Do not use more frequently than once every week.

PREMISE SPRAY FOR YARDS, KENNELS: Mix at the rate of 2 1/2 fl. ozs. (5 tablespoonfuls) with 1 gallon of water. Use 1 gallon of spray per 1000 sq. ft. of lawn or kennel area. Make sure to spray possible hiding places such as cracks in cement, under and around dogs' bedding and under various pieces of debris.

CATTLE, SHEEP, GOATS, SWINE, AND HORSES

TO KILL TICKS, HORN FLIES, LICE, KEDS, WOOL MAGGOTS AND SCREWORMS: Mix at the rate of 1 gallon Co Nav in 200 gallons of water. Mix thoroughly. Spray or dip as required but no more often than every two weeks. The use of Co Nav requires no waiting period between last application and slaughter.

LONG-LASTING PROTECTION: Co Nav provides long-lasting protection against reinfestation of ticks and 3-week protection against horn flies. A single application is usually sufficient for lice and keds. For biting lice on goats, a second application after 21 days may be required for complete clean-up. Co Nav also aids in controlling screwworm larvae and provides up to 2 months protection against wool maggots.

SPRAYING: Make sure spray is thoroughly mixed before application. Wet animals thoroughly for best results. Do not allow animals to drink from run-off pools.

DIPPING: Thoroughly mix dip before dipping. Water animals before dipping to prevent drinking of dip wash. In routine dipping, if it is necessary to treat animals under the age of 3 months, spraying is recommended. In quarantine dipping for fever tick control, when animals of all ages must be dipped, individual hand immersion of young animals is essential to avoid swallowing of dip wash.

REPLENISHMENT: Add 1 gallon of Co Nav to each full gallon of fresh water added to dipping vat or control replenishment by vat-side test.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinse liquids that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Triple rinse or equivalent and dispose of in an approved landfill or bury in a safe place.

GENERAL: Consult Federal, State or local disposal authorities for approved alternative procedures.

WELLCOME VETERINARY DIVISION
Burroughs Wellcome Co.
Research Triangle Park, N.C. 27709

59-113