

## Co-Nav

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (& DOMESTIC ANIMALS) WARNING

Co-Nav is toxic and may be fatal if swallowed or absorbed through the skin. Do not inhale spray, get in eyes, on skin or on clothing. Use respirator or goggles if necessary. Wash with soap and water in case of contact. Do not use on dairy animals or in dairy barns. Do not dip or spray foals. Do not dip animals under 3 months of age.

Do not contaminate feed or foodstuffs. Food utensils such as teaspoons and tablespoons should not be used for food purposes after use with pesticides.

Apply this product only as specified on this label.

### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician, veterinarian or Poison Control Center immediately. DO NOT INDUCE VOMITING, unless under medical supervision. Vomiting may cause aspiration pneumonia.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

IF ON SKIN: Wash immediately with plenty of soap and water.

IF IN EYES: Flush eyes with plenty of water. See physician immediately.

NOTE TO PHYSICIAN, VETERINARIAN: This product contains an organophosphate and petroleum distillates. ATROPINE IS ANTIDOTAL for the organophosphate ingredient.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish, wildlife and birds. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where run-off is likely to occur.

### PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**DOGS, TICKS, FLEAS AND LICE:** Thoroughly mix at the rate of 1/2 fl. oz (1 tablespoonful) with 1 gallon of water.

Sponge, swab, bathe or dip dog making sure dog is thoroughly wet.

Repeat only when necessary but not more frequently than once every week.

**PREMISE SPRAY FOR YARDS, KENNELS:** Mix at the rate of 2 1/2 fl. ozs (5 tablespoonful) Use 1 gallon of spray per 1000 sq. ft of lawn or kennel area. Make sure to spray possible cracks in cement or wood, around dog's bedding and under various pieces of debris.

### CATTLE, SHEEP, GOATS, SWINE, AND HORSES

**TO KILL TICKS, HORN FLIES, LICE, KEDS, WOOL MAGGOTS AND SCREW-WORMS:** Mix at 1 in 200 gallons of water. Mix thoroughly. Spray or dip as required but no more often than of Co-Nav requires no waiting period between last application and slaughter.

**LONG-LASTING PROTECTION:** Co-Nav provides long-lasting protection against reinfestation against horn flies. A single application is usually sufficient for lice and keds. For second application after 21 days may be required for complete clean-up. Co-Nav also aids larvae and provides up to 2 months protection against wool maggots.

**SPRAYING:** Make sure spray is thoroughly mixed before application. Wet animals thoroughly allow animals to drink from run-off pools.

**DIPPING:** Thoroughly mix dip before dipping. Water animals before dipping to prevent dry time dipping, if it is necessary to treat animals under the age of 3 months, spraying is recommended for fever tick control when animals of all ages must be dipped, individual handling is essential to avoid swallowing of dip wash.

**REPLENISHMENT:** Add 1 gallon of Co-Nav to each 200 gallons fresh water added to dipping ment by vat-side test.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in cool, dry place away from heat or open flame.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture or rinse liquids that cannot be used, instructions must be disposed of according to Federal, State or local procedures under Pollution Control Act.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent) and dispose of in a sanitary landfill. State and local procedures.

WELLCOME ANIMAL HEALTH DIVISION

Burroughs Wellcome Co.

Kansas City, MO 64108

Made in U.S.A.

**CAUTIONARY STATEMENTS  
FOR HUMANS (& DOMESTIC ANIMALS)**

**WARNING**  
Avoid contact with or absorbed through the skin. Do not inhale spray, get in eyes. Wash with soap and water in case of contact. Do not dip or spray foals. Do not dip animals under 3 months of age.

Food utensils such as teaspoons and tablespoons should not be used with this product. See label.

**SYMPTOMS OF PRACTICAL TREATMENT**

See physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia.

Apply respiration if indicated.

Wash with soap and water.

See physician immediately.

This product contains an organophosphate and petroleum distillates. See label for phosphorus ingredient.

**ENVIRONMENTAL HAZARDS**

Keep out of any body of water. Do not contaminate water by runoff. Do not apply where runoff is likely to occur.

**PHYSICAL OR CHEMICAL HAZARDS**

See label.

**DIRECTIONS FOR USE**

Use in a manner inconsistent with its labeling.

Mix at the rate of 1/2 fl oz (1 tablespoonful) with 1 gallon of water. Spray or dip thoroughly wet.

Repeat frequently than once every week.

**PREMISE SPRAY FOR YARDS, KENNELS:** Mix at the rate of 2 1/2 fl. ozs (5 tablespoonfuls) with 1 gallon of water. Use 1 gallon of spray per 1000 sq. ft of lawn or kennel area. Make sure to spray possible hiding places such as cracks in cement or wood, around dog's bedding and under various pieces of debris.

**CATTLE, SHEEP, GOATS, SWINE, AND HORSES**

**TO KILL TICKS, HORN FLIES, LICE, KEDS, WOOL MAGGOTS AND SCREWORMS:** Mix at the rate of 1 gallon Co Nav in 200 gallons of water. Mix thoroughly. Spray or dip as required but no more often than every two weeks. The use of Co Nav requires no waiting period between last application and slaughter.

**LONG-LASTING PROTECTION:** Co Nav provides long-lasting protection against reinfestation of ticks and 3-week protection against horn flies. A single application is usually sufficient for lice and keds. For biting lice on goats, a second application after 21 days may be required for complete clean-up. Co Nav also aids in controlling screwworm larvae and provides up to 2 months protection against wool maggots.

**SPRAYING:** Make sure spray is thoroughly mixed before application. Wet animals thoroughly for best results. Do not allow animals to drink from run-off pools.

**DIPPING:** Thoroughly mix dip before dipping. Water animals before dipping to prevent drinking of dip wash. In routine dipping, if it is necessary to treat animals under the age of 3 months, spraying is recommended. In quarantine dipping for fever tick control when animals of all ages must be dipped, individual hand immersion of young animals is essential to avoid swallowing of dip wash.

**REPLENISHMENT:** Add 1 gallon of Co Nav to each 200 gallons fresh water added to dipping vat or control replenishment by vat-side test.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in cool, dry place away from heat or open flame.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture or rinse liquids that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent) and dispose of in a sanitary landfill or by other approved State and local procedures.

**WELLCOME ANIMAL HEALTH DIVISION**

Burroughs Wellcome Co.

Kansas City, MO 64108

Made in U.S.A.

112402