

# Hog Mange Cure

## For Sarcoptic Mange, Kills Lice

### Active Ingredients

Toxaphene (Technical Chlorinated Camphene containing 67 to 69% Chlorine).....	45.00%
Gamma Isomer of Benzene Hexachloride (from lindane).....	1.95%
Xylene.....	5.01%
Kerosene.....	34.94%
Inert Ingredients.....	13.09%
	100.00%

**WARNING:** Keep out of reach of children. See other warnings on back panel.

Net Contents One Gallon (U.S. Measure)

EPA Reg. No. 59-127-AA



## Hog Mange

### DIRECTIONS FOR

**Sarcoptic Mange**  
Mix Cooper Hog Mange  
twice at a 14 day inter

**Lice**  
Mix Cooper Hog Mange  
application is usually s

**Hog Rubbers**  
To aid in controlling lice  
1 1/2 gallons Diesel fuel  
so it is thoroughly wet,  
necessary. Regular exp  
good results against lice

Amount of Cooper Hog Mange Cure
1/2 pint
1 pint
1 quart
1 gallon

# Hog Mange Cure

## DIRECTIONS FOR USE

### Sarcoptic Mange

Mix Cooper Hog Mange Cure at the rate of 1 pint to 12 gallons of water. Spray hogs twice at a 14 day interval. Thorough wetting of animals is essential.

### Lice

Mix Cooper Hog Mange Cure at the rate of 1 pint to 20 gallons of water. A single application is usually sufficient for control of this pest.

### Hog Rubbers

To aid in controlling lice, mix Cooper Hog Mange Cure at the rate of one pint to 1½ gallons Diesel fuel or light motor oil. Charge burlap roll by soaking with solution so it is thoroughly wet, but not to the point of dripping. Repeat charging as necessary. Regular exposure of animals for about 10 weeks is usually necessary for good results against lice.

### Dilution Table

Amount of Cooper Hog Mange Cure	Amount of water to use		Hog Rubbers
	Sarcoptic Mange	Lice	
½ pint	6 gal.	10 gal.	3 qts. oil
1 pint	12 gal.	20 gal.	1½ gal. oil
1 quart	25 gal.	40 gal.	3 gal. oil
1 gallon	100 gal.	160 gal.	12 gal. oil

## WARNING

May be fatal if swallowed. This product is toxic and is absorbed through the skin in dry form and from solutions. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Use respirator or goggles if necessary. Do not contaminate food or food products. Concentrations of Cooper Hog Mange Cure greater than those recommended should never be used. Do not use on household pets or humans.

Do not treat animals under 3 months of age. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter. Avoid treating animals in cold stormy weather. Avoid treating overheated or sick animals. Do not expose animals under 3 months of age to backrubber. Do not use on animals within 45 days of slaughter.

This product is toxic to fish, wildlife, and birds. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

### Antidote

Human—Internal: Call a doctor immediately. Give sodium chloride 15 gms. (one tablespoonful of salt) in a glass of warm water. Repeat until vomit is clear.  
External: Wash with plenty of soap and water.

## COOPER U.S.A., INC.

Research Triangle Park, N.C. 27709

A Wellcome Company

EPA Est. 59-NC-1

Q25600