

Prolin* Rat & Mouse Killer "Feeder Pak"

DIRECTIONS FOR USE

Rats

Place a minimum of 1 packet (4 ounces) of bait per station in dry containers next to burrows, in runways or in locations where rats feed, water or travel. Bait placements along walls, in corners, and in concealed locations where rats can feed unobserved are desirable. Replace contaminated, stale, damp or moldy baits with fresh bait promptly. Assure an uninterrupted supply of bait for at least 15 days and continue baiting until all signs of feeding have stopped. Where a continuous source of infestation is present (nearby dumps) permanent bait stations should be established and replenished as needed. Each 4 oz. feeder pak contains enough bait to kill 1 or 2 rats.

Mice

Place tablespoon amounts (¼ to ½ ounce) of bait in shallow containers where mice or signs of mice have been seen and where mice feed, water, or travel. Bait stations at intervals of 8 to 12 feet are recommended. Bait placements along walls, in corners, and in concealed locations where mice can feed unobserved are desirable. Replace contaminated, stale, damp, or moldy baits with fresh bait promptly. Assure an uninterrupted supply of bait for at least 15 days and continue baiting until all signs of feeding have stopped.

Prolin* Rat & Mouse Killer "Feeder Pak"

Warfarin plus Sulfaquinoxaline Rodenticide
Ready-to-Use "Pelletized" Bait
Packed in 72 x 4 oz. "Feeder Paks" for Easy Baiting

Active Ingredients

N'-(2-quinoxaliny) sulfanilamide (Sulfaquinoxaline)..... 0.025%
Warfarin 3-(α -acetonylbenzyl)-4-hydroxycoumarin..... 0.025%

Inert Ingredients..... 99.950%
..... 100.000%

CAUTION: Keep out of reach of children.

See other cautions on side panel.


*TM Wisconsin Alumni Research Foundation

Net Contents 18 Pounds

EPA Reg. No. 59-137



COOPER



Prolin* Rat & Mouse Killer "Feeder Pak"

CAUTION

Protect children, pets, wildlife and domestic animals from bait. Bait should be placed in locations inaccessible to children, pets, wildlife and domestic animals or in tamper proof boxes. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains warfarin; an anticoagulant chemical that reduces the clotting ability of the blood and upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains sulfaquinoxaline; an antibacterial agent which inhibits bacteria that produce vitamin K, an antidote for warfarin.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

Manufactured For
COOPER U.S.A., INC.
Chicago, Illinois 60614, USA

A Wellcome Company

VWSEW-L-MWWH

COOPER PROLIN®
RAT & MOUSE KILLER
FEEDER-PAK

©Prolin is a registered trademark of the
Wisconsin Alumni Research Foundation.

DIRECTIONS FOR USE

RATS—Place a minimum of 1 packet (4 ounces) of bait per station in dry containers next to burrows, in runways or in locations where rats feed, water, or travel. Bait placements along walls, in corners, and in concealed locations where rats can feed unobserved are desirable. Replace contaminated, stale, damp or moldy baits with fresh bait promptly. Assure an uninterrupted supply of bait for at least 15 days and continue baiting until all signs of feeding have stopped. Where a continuous source of infestation is present (nearby dumps) permanent bait stations should be established and replenished as needed. This Feeder Pak contains enough bait to kill 1 or 2 rats.

MICE—Place tablespoon amounts (1/4 to 1/2 ounce) of bait in shallow containers where mice or signs of mice have been seen and where mice feed, water, or travel. Bait stations at intervals of 8 to 12 feet are recommended. Bait placements along walls, in corners, and in concealed locations where mice can feed unobserved are desirable. Replace contaminated, stale, damp or moldy baits with fresh bait promptly. Assure an uninterrupted supply of bait for at least 15 days and continue baiting until all signs of feeding have stopped.

CAUTION: Keep out of reach of children.
See other cautions on front panel.

Net Weight 4 Oz.

