

Cio-Rid Ciodrin* Emulsifiable Liquid

DIRECTIONS FOR USE

Dairy Cattle

To control Face Flies, Stable Flies, Horn Flies, Ticks and Lice. Dilute 1 pint of Cio-Rid in 3½ gallons of water (1 to 28). Apply one to two quarts of dilute spray per adult animal and proportionately less for younger animals. Repeat as required but not more often than once every seven days.

Loose Penned and Feed Lot Cattle

To control Face Flies, Stable Flies, Horn Flies and Lice. Dilute 1 pint of Cio-Rid in 3½ gallons of water (1 to 28). Spray at the rate of 1 to 2 quarts of diluted spray over and on adult animals and proportionately less for younger animals. Repeat as required but not more often than once every seven days.

Sheep, Goats and Swine

To control Stable Flies, Horn Flies, Face Flies, Lice on Swine, Lice and Ticks on Sheep and Goats. Dilute 1 pint of Cio-Rid in 6 gallons of water (1 to 50) and spray at the rate of one gallon of diluted spray per animal. Repeat as required but not more often than once every seven days.

Back Rubbers

For control of Flies on Beef and Dairy Cattle. Dilute 1 pint of Cio-Rid with 2¼ gallons of suitable oil. Charge back rubber as needed. Do not spray animals with this oil solution.

Surface Spray

For control of House Fly and Resting Stable Fly. Dilute 1 pint of Cio-Rid in 1¼ gallons of water (1 to 14). Spray at the rate of 1 gallon of diluted spray per 1000 square feet of surface (wet surfaces, but not to point of excessive run-off). Spray walls of dairy barns, on fences, hog pens and other similar places where flies are seen to congregate. Do not apply Ciodrin* (Cio-Rid) in poultry houses. Repeat as required or at least twice monthly.

Spray Dilution Rates

(1 to 28)		(1 to 50)	
Cio-Rid	Water	Cio-Rid	Water
8 Fluid Ounces (1 measuring cup)	1¾ Gallons	5 Tablespoons	1 Gallon
1 Pint	3½ Gallons	8 Fluid Ounces (1 measuring Cup)	3 Gallons
1 Quart	7 Gallons	1 Pint	6 Gallons
1 Gallon	28 Gallons	1 Quart	12 Gallons
		1 Gallon	48 Gallons

WARNING

HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN.

Avoid contact with skin, eyes, or clothing. Do not breathe spray mist. During commercial exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type found suitable by the U. S. Bureau of Mines for protection against Ciodrin* insecticide. Apply this product only as specified on this label and at recommended dosages. Wash thoroughly with soap and water after handling and before eating and smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Keep away from heat and open flame. Keep out of reach of children.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not contaminate feed or foodstuffs. Food utensils such as teaspoons and tablespoons should not be used for food purposes after use with pesticides.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

COOPER U.S.A., INC.

Chicago, Illinois 60614, USA

A Wellcome Company

Cooper-Tox Extra Dust

Kills Horn Flies, Lice and Ticks on Beef Cattle, Sheep, Goats and Swine

Active Ingredients

Permethrin (0.5% active ingredient)	5
Pyrethrin (0.1% active ingredient)	1

Inert Ingredients

94
100%

CAUTION: Keep out of reach of children. See other cautions below.

DIRECTIONS FOR USE

Beef Cattle - Apply to the head, neck, and body of the animal. Use a spray nozzle or dust applicator. Repeat every 7 to 10 days.

Sheep and Goats - Apply to the head, neck, and body of the animal. Use a spray nozzle or dust applicator. Repeat every 7 to 10 days.

Swine - Apply to the head, neck, and body of the animal. Use a spray nozzle or dust applicator. Repeat every 7 to 10 days.

Precautions - Do not use on animals that are pregnant or nursing. Do not use on animals that are sick or stressed. Do not use on animals that are being transported.

CAUTION

Keep out of reach of children. Do not use on animals that are pregnant or nursing. Do not use on animals that are sick or stressed. Do not use on animals that are being transported. Do not use on animals that are being sold or shown. Do not use on animals that are being used for food. Do not use on animals that are being used for wool. Do not use on animals that are being used for skins. Do not use on animals that are being used for hides. Do not use on animals that are being used for bones. Do not use on animals that are being used for horns. Do not use on animals that are being used for hooves. Do not use on animals that are being used for hoofs. Do not use on animals that are being used for hooves. Do not use on animals that are being used for hooves.

Net Weight 5 Pounds

Cooper U.S.A., Inc.
Storage: 50-104-007
A Wellcome Company

EPA Reg. No. 53-100

