

[59-155]

Caution.

- Harmful if swallowed, inhaled or absorbed through the skin.
- Do not breathe spray, mist or vapor.
- Do not get on skin.
- Wash with soap and water after using and before eating or smoking.
- Wash clothing before re-use.
- Do not contaminate feed or foodstuffs.
- Keep out of reach of children and domestic animals.

This product is toxic to fish. Keep out of lakes, ponds, streams, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Disposal of Containers:

Do not reuse empty drum. Return to drum in condition or destroy by perforating or crushing and burying in a safe place.

RECEIVED

SEP 24 1976

UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND ROYACIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 27272 SUBJECT TO ATTACHED CONDITIONS

COOPER MALATHION EM #5

ACTIVE INGREDIENTS

Malathion (C,2 dimethyl phosphorodithioate of diethyl mercaptosuccinate) 5.49%
Aromatic Petroleum Solvent 45.89%
INERT INGREDIENTS 48.62%

*One gallon contains 5 pounds of Malathion
E.P.A. Reg. No. 59-155

CAUTION:

Keep out of reach of children.
See other cautions on back panel.

For control of Lice, Ticks & Larvae on Bees, Cattle and Horses, Lice and Sarcoptic Mite on Hogs, Lice, Ticks & Keds on Sheep and Goats, Flies and Mosquito larvae.

William Cooper & Nephews, Inc.

1909 North Clinton Avenue

Chicago, Illinois 60641

Telephone: 312-349-6000

Livestock: Use the following

Pest	Amount of Malathion
Bees	5 1/2 gal
Cattle (also not lactating dairy animals)	5 gal
Horses	5 gal
Hogs	5 gal
Sheep and Goats	5 gal

Do not apply to lactating dairy and breeding. Do not treat animals one hour before or after feeding. Do not contaminate feed or food containers.

Check: Use the following table for the pests listed below.

Pest	Amount Malathion EM-50	Remarks
Lice	4 1/2 oz in 5 gal water	Thorough wetting of animals is essential for best results. Repeat application at 10 days and thereafter as needed.
Ticks	6 1/2 oz in 5 gal water	Thorough wetting of animals is essential for best results. Repeat application at 2 week intervals as needed.
Horn flies	6 1/2 oz in 5 gal water	Thorough wetting of animals is essential for best results. Repeat application at 2 week intervals as needed.
Lice	4 1/2 oz in 5 gal water	Thorough wetting of animals is essential for best results. Apply a complete coverage spray to animals' heads and other. Repeat at 10 day intervals as needed.
Sarcoptes Mange	4 1/2 oz in 5 gal water	Apply to animals' heads and other. Repeat at 10 day intervals as needed.
Lice	4 1/2 oz in 5 gal water	Thorough wetting of animals is essential for best results. Apply a complete coverage spray to animals' heads and other. Repeat at 10 day intervals as needed.
Ticks	6 1/2 oz in 5 gal water	Thorough wetting of animals is essential for best results. Repeat application at 2 week intervals as needed.
Kills	4 1/2 oz in 5 gal water	Thorough wetting of animals is essential for best results. Repeat application at 2 week intervals as needed.

Do not apply to lactating dairy animals or nonlactating dairy animals within 2 weeks of milking. **Do not** treat animals under 1 month of age. When applying sprays, avoid contamination of feed, food containers and watering troughs.

Check: Use the following table for the pests listed below.

Pest	Amount Spray	Amount Malathion EM-50	Remarks
Lice	4 1/2 Gals	4 1/2 Gals	Thorough wetting of animals is essential for best results. Repeat application at 10 days and thereafter as needed.
Ticks	12 Gals	12 Gals	Thorough wetting of animals is essential for best results. Repeat application at 2 week intervals as needed.

Apply as a spray at the rate of 1/2 ounce of active ingredient per 1000 gallons of water. Apply to the body of the animal, avoiding the face and eyes. Thorough wetting of animals is essential for best results. Repeat application at 10 day intervals as needed. Avoid contamination of feed, food containers and watering troughs. Remove lactating dairy animals from milking areas before treatment. Also, do not milk animals under 1 month of age before treatment.

Mosquito Control:

Apply to water in watering troughs, feed buckets, and other containers. Avoid contamination of feed, food containers and watering troughs. Remove lactating dairy animals from milking areas before treatment. Also, do not milk animals under 1 month of age before treatment.