

**Container Disposal:** Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Then dispose of bag in a sanitary landfill or by incineration if allowed by State and local authorities.

**Mellcor Animal Health, Inc.**  
Kansas City, MO 64108 U.S.A.

**To drop insecticide into burlap  
area, pull zipstring.**

Wellcome Animal Division  
Swaroughs Wellcome Co.  
2000 South 11th St.  
Kansas City, KS 64103

300 / 116869  
17 / 3

Attention: Donald A. Hunt

Gentlemen:

Subject: Amendment - Updated Label  
Cooper Cattle Dust No. C  
EPA Registration No. 59-156  
Application Dated October 14, 1983

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Fodenticide Act (FIFRA), is acceptable provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

1. Based upon a request from Bayvet to reconsider the toxicity category for 1% and less coumaphos dust formulations, the Agency has decided to go with a toxicity category III classification for these products and therefore is requiring the signal word "CAUTION." You should make this revision to your labeling.
2. Also delete "May be fatal if inhaled," and revise "Harmful if swallowed or absorbed through skin" to read "Harmful if swallowed, inhaled, or absorbed through skin."

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George S. FERGUSON  
Product Manager (15)  
Insecticide Registration Branch  
Registration Division (21-767)

Enclosure

10-21-83 1-95517:1 (2) 20:1 10:1 -17:4730 00:01:5 13/13

CONFERENCE							
SYMBOL							
SURNAME	T. Hunt						
DATE							

Bag - Cattle Dust Bag C  
(Front Panel)

(YELLOW BACKGROUND)

Cattle Dust Bag C

For Control of Lice and Horn Flies and as an Aid in Reducing Face Fly  
Population on Beef and Dairy Cattle

Active Ingredient

0,0-Diethyl O-(3-chloro-4-methyl-2-oxo-(2H)-

1-benzopyran-7 yl) phosphorothioate..... 1%

Inert Ingredients..... 99%

100%


KEEP OUT OF REACH OF CHILDREN  
WARNING  
SEE BACK PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS

ACCEPTED  
with COMMENTS  
MAY 1964  
59-156

Net Weight 25 Pounds

EPA Est. 59-NC-1

EPA Reg. No. 59-156



(RED CIRCLE  
WHITE REVERSE UNICORN  
BLACK BACKGROUND)

(WHITE REVERSE TYPE)

(BLACK BAR)

(Back Panel)

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
WARNING

Harmful if swallowed or absorbed through skin. May be fatal if inhaled. Do not breathe dust. Do not get in eyes, on skin or on clothing. Wash hands thoroughly after handling product and before eating or smoking.

STATEMENT OF PRACTICAL TREATMENT

**If Swallowed:** Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available, by administering syrup of ipecac. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

**If Inhaled:** Remove victim to fresh air and apply respiration if indicated.

**If On Skin:** Remove contaminated clothing and immediately wash skin thoroughly with soap and water.

**If In Eyes:** Immediately flush eyes thoroughly for at least 5 minutes with plenty of water. Get medical attention if irritation persists.

**Note To Physician:** Prolonged exposure will result in cholinesterase depression. ATROPINE IS ANTIDOTAL. 2-PAM is also antidotal and may be administered in conjunction with atropine.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of lakes streams or ponds. Do not contaminate water by cleaning of equipment or by disposal of wastes. Apply this product only as specified on this label.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Important:** Read entire label carefully before using Cattle Dust Bag C. Use only as directed.

**Louse and Horn Fly Control:** Suspend Cattle Dust bag C in areas frequented by cattle or in gateways or lanes through which the animals pass daily for water, feed or minerals. Bags may also be suspended in loafing sheds or in front of protected mineral feeders. For lactating dairy cows, Cattle Dust Bag C may be suspended in the exit through which cows leave the milking barn. In all cases, the dust bags should be adjusted so that the bottom of the bag will hang 4 to 6 inches below the top line of the cattle.

**Louse and Horn Fly Control and as an Aid in Reducing Face Fly Populations:** Suspend Cattle Dust Bag C in the same manner as described under Louse and Horn Fly Control except as follows: (1) The bottom of the bags should be 18 to 24 inches from the ground to insure that the faces of the cattle will be dusted. (2) The bags must be located so the animals will be forced to use them daily.

As face flies are able to travel over an area of several miles, best results in reducing face fly populations are achieved when the dust bags are used on an area-wide basis.

**Installation of Cattle Dust Bag C:** Suspend the dust bag by inserting rope, chain or wire through the grommets at the top of the bag and attaching to an overhead structure or by hooking the grommets into hooks fixed to a stationary overhead 2 x 4 at the height recommended for louse and horn fly or face fly control. In areas of frequent or heavy rainfalls, Cattle Dust Bag C should be protected from the weather.

Install as many dust bags as necessary in lanes or other openings to assure proper dusting of the animals. After the bags are in place, pull the rip cord to allow the insecticide to fall into the bottom of bag. When dust bags are provided on a free choice basis, allow one 25 pound Cattle Dust Bag C per 25 to 50 animals.

**Refilling Cattle Dust Bag C:** Starting at top of bag near grommets, slit either side seam approximately 10 inches downward from bottom of grommets. This will make an opening large enough to insert the unopened refill cartridge partially into the bag. Slit open the refill cartridge inside the dust bag, and empty entire contents. It is not necessary to take the dust bag from its hanging position to complete this operation.

**Use Restrictions:** For external insecticidal use on specified animals only. Do not contaminate feed, troughs, water or water utensils. Provide thorough ventilation while dusting. Do not apply to sick, stressed or convalescent animals.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Storage:** Store in cool, dry place.

**Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.