

Directions For Use

Important: Read entire label carefully before using CATTLE DUST BAG C. Use only as directed.

Louse and Horn Fly Control: Suspend Cattle Dust Bag C in areas frequented by cattle or in gateways or lanes through which the animals pass daily for water, feed or mineral. Bags may also be suspended in loafing sheds or in front of protected mineral feeders. For lactating dairy cows Cattle Dust Bag C may be suspended in the exit through which cows leave the milking barn. In all cases, the dust bags should be adjusted so that the bottom of the bag will hang four to six inches below the topline of the cattle.

Louse and Horn Fly Control and as an aid in reducing Face Fly populations: Suspend Cattle Dust Bag C in the same manner as described under Louse and Horn Fly Control except as follows: (1) The bottom of the bags should be 18 to 24 inches from the ground to insure that the faces of the cattle will be dusted. (2) The bags must be located so the animals will be forced to use them daily.

As face flies are able to travel over an area of several miles, best results in reducing face fly populations are achieved when the dust bags are used on an area-wide basis.

Installation of Cattle Dust Bag C: Suspend the dust bag by inserting rope, chain, or wire through the grommets at the top of the bag and attaching to an overhead structure or by hooking the grommets into hooks fixed to a stationary overhead 2 x 4 at the height recommended for louse and horn fly or face fly control. In areas of frequent or heavy rainfall, Cattle Dust Bag C should be protected from the weather.

Install as many dust bags as necessary in lanes or other openings to assure proper dusting of the animals. After the bags are in place, pull the rip cord to allow the insecticide to fall into the bottom of the bag. When dust bags are provided on a free-choice basis, allow one 25-pound Cattle Dust Bag C per 25 to 50 animals.

Refilling Cattle Dust Bag C: Starting at top of bag near grommets, slit either side seam approximately 10 inches downward from bottom of grommets. This will make an opening large enough to insert the unopened refill cartridge partially into the bag. Slit open the refill cartridge inside the dust bag, and empty entire contents. It is not necessary to take the dust bag from its hanging position to complete this operation.

Do not reuse refill cartridge bag. Destroy when empty.

Restrictions: Do not hang dust bag over feed, mineral or water troughs.

Caution: Harmful if inhaled or absorbed through the skin. Do not get in eyes or on skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling.

Do not contaminate food or feed. Keep out of reach of children.

Destroy damaged or worn-out dust bag by burying or burning. Stay away from smoke or fumes.

This product is toxic to fish, birds and other wildlife. Do not contaminate water by cleaning of equipment or by disposal of wastes. Apply this product only as specified on this label.

If poisoning occurs obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

To Physician: Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

Cooper U.S.A., Inc.
Chicago, Illinois 60614 USA
A Wellcome Company

To drop insecticide into burlap area, pull zipstring.