



Bag - Dust-Pak
(Front Panel)

(YELLOW BACKGROUND)

Dust-Pak
Cattle Dust Bag Filler

For Control of Lice and Horn Flies and as an Aid in Reducing Face Fly
Population on Beef and Dairy Cattle

Active Ingredient

0,0-Diethyl 0-(3-chloro-4-methyl-2-oxo (2H)-

1-benzopyran-7 yl) phosphorothioate..... 1%

Inert Ingredients..... 99%

100%

KEEP OUT OF REACH OF CHILDREN
WARNING
SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

ACCEPTED
with COMMENTS
by EPA
MAY 15 1964
59-174

Net Weight 10 Pounds

EPA Est. 59-NC-1

EPA Reg. No. 59-174

 **COOPER**

(RED CIRCLE
WHITE REVERSE UNICORN
BLACK BACKGROUND)

(WHITE REVERSE TYPE)

(BLACK BAR)

(Back Panel)

Dust-Pak Cattle Dust Bag Filler

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING**

Harmful if swallowed or absorbed through skin. May be fatal if inhaled. Do not breathe dust. Do not get in eyes, on skin or on clothing. Wash hands thoroughly after handling product and before eating or smoking.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available, by administering syrup of ipecac. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If Inhaled: Remove victim to fresh air and apply respiration if indicated.

If On Skin: Remove contaminated clothing and immediately wash skin thoroughly with soap and water.

If In Eyes: Immediately flush eyes thoroughly for at least 5 minutes with plenty of water. Get medical attention if irritation persists.

Note To Physician: Prolonged exposure will result in cholinesterase depression. ATROPINE IS ANTIDOTAL. 2-PAM is also antidotal and may be administered in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of lakes streams or ponds. Do not contaminate water by cleaning of equipment or by disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Filling Dust Bags with Dust-Pak

Open dust bag and partially insert the unopened Dust-Pak. Slit open Dust-Pak inside the bag, and empty entire contents. Cooper dust bags will hold two Dust-Paks.

Installation of Dust Bag

Suspend dust bag by inserting rope, chain or wire through grommets at top of bag and attaching to an overhead structure or by hooking grommets into hooks fixed to a stationary overhead 2 x 4 at the height recommended for louse and horn fly or face fly control. In areas of frequent or heavy rainfall, dust bags should be protected from the weather.

Install as many dust bags as necessary in lanes or other openings to assure proper dusting of the animals. When dust bags are provided on a free choice basis, allow one dust bag for 25 to 50 animals.

Louse and Fly Control with Dust Bags

Dust-Pak is designed for use with Cooper Cattle Dust Bags, but may be used as a refill for any standard dust bag.

Important: Read entire label carefully before using Dust-Pak. Use only as directed.

Louse and Horn Fly Control

Suspend dust bags in areas frequented by cattle or in gateways or lanes through which the animals pass daily for water, feed or minerals. Bags may also be suspended in loafing sheds or in front of protected mineral feeders. For lactating dairy cows, dust bags may be suspended in the exit through which cows leave the milking barn. In all cases, the dust bags should be adjusted so that the bottom of the bag will hang 4 to 6 inches below the top line of the cattle.

Louse and Horn Fly Control and as an Aid in Reducing Face Fly Populations

Suspend dust bags in the same manner as described under Louse and Horn Fly Control but the bottom of the bags should be low enough to insure that the faces of the cattle will be dusted. The bags must be located so that the animals will be forced to use them daily.

As face flies are able to travel over an area of several miles, best results in reducing face fly populations are achieved when the dust bags are used on an area-wide basis.

Use Restrictions

For external insecticidal use on specified animals only. Do not contaminate feed, troughs, water or water utensils. Provide thorough ventilation while dusting. Do not apply to sick, stressed or convalescent animals.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in cool, dry place.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Wellcome Animal Health, Inc.
Kansas City, MO 64108 U.S.A.

