



For Control of Lice and Horn Flies on Beef and Dairy Cattle and Lice on Swine

Active Ingredient

0,0-Diethyl 0-(3-chloro-4-methyl-2-oxo-(2H)-1-benzopyran-7-yl) phosphorothioate 1%

Inert Ingredients 99%

100%

KEEP OUT OF REACH OF CHILDREN
CAUTION
 SEE BACK PANEL FOR ADDITIONAL
 PRECAUTIONARY STATEMENTS

526163

Net Weight 25 Pounds (11.3 kg)

EPA Est. 602-MO-1
EPA Reg. No. 59-175



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Do not breathe dust. Do not get in eyes, on skin, or on clothing. Wash hands thoroughly after handling product and before eating or smoking.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available, by administering syrup of ipecac. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If Inhaled: Remove victim to fresh air and apply respiration if indicated.

If On Skin: Remove contaminated clothing and immediately wash skin thoroughly with soap and water.

If In Eyes: Immediately flush eyes thoroughly for at least 5 minutes with plenty of water. Get medical attention if irritation persists.

Note To Physician: Prolonged exposure will result in cholinesterase depression. ATROPINE IS ANTIDOTAL. 2-PAM is also antidotal and may be administered in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or by disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Important: Read the entire label carefully before using Dri-Kil Extra. Hold container at a distance from the area to be treated that will permit even distribution of the dust. 1 oz of Dri-Kil Extra equals approximately three level tablespoonsful.

Beef and Dairy Cattle: To Control Lice and Horn Flies: Apply not more than 2 oz Dri-Kil Extra dust per animal. Dust evenly into hair over the head, neck, shoulders, back and tailhead. Repeat as necessary. No interval is required between treatment and use of meat or milk for food.

Swine: To Control Lice: Apply not more than 1 oz of Dri-Kil Extra dust per animal as a uniform coat over the sides and back. Repeat as necessary but not more often than once every 10 days. No interval is required between treatment and use of meat as food.

Bedding: Apply 2 oz of Dri-Kil Extra dust evenly over each 30 square feet of fresh, dry bedding. Repeat as necessary but not more often than once every 10 days.

Note: In severe infestations, both individual animal and bedding may be treated as directed above.

Use Restrictions: For external insecticidal use on the above specified animals only. Do not contaminate feed, troughs, water or water utensils. Provide thorough ventilation while dusting. Do not apply to sick, stressed or convalescent animals.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in cool, dry place.

Pesticide Disposal: Pesticides wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Wellcome Animal Health, Inc.
Kansas City, MO 64108 U.S.A.
12/26/83