

Reg # 59-201

PM-18

191

NOTIFICATION
LABEL NOT REVENUE
PER FR NOTICE 06-3
DATE ~~MAY 9~~ 1991

Salsbury™ Permaban® 11% EC

Emulsifiable concentrate long-lasting
insecticidal spray for livestock and
their premises

EPA Est 6175 LA 1
EPA Reg No 59 201 116

Active ingredient
Permethrin (3-phenoxyphenyl) methyl (±) cis,
trans-3-(2,2-dichloroethyl)-2,2-dimethyl-
cyclopropanecarboxylate^a 11.0%
Inert ingredients 89.0%
100.0%

^aActive ingredient: 100% (±) cis and 100% (±) trans
This product contains 8.187 lbs. (48.5 g)
of permethrin per pint (473 ml)

Keep out of reach of children
Caution
See side panel for additional precautionary
statements

Net Contents: One U.S. Pint (473 ml)

SOLVAY Animal Health, Inc.

Directions for use
It is a violation of Federal law to use this product in a manner inconsistent with its
labeling
Mix Salsbury Permaban 11% EC thoroughly according to the following table:
Repeat applications as needed, but not more often than once every two weeks. Spray
lactating dairy animals only after milking is completed. Use as needed on horses not
intended for human consumption.

Target species	Use	Dilute	Application Rate
Lactating and non-lactating dairy cattle and goats, beef cattle, horses and sheep			
Horn flies only	Sprayer	1 pt to 50 gals water (3 tbsp/5 gals)	1 qt of coarse spray per animal
Horn flies, Face flies, Stable flies, House flies, Horse flies, Black flies, Mosquitoes, Eye gnats, Mange mites, Scabies mites, Ticks, Lice, Sheep keds	Sprayer	1 pt to 25 gals water (3 tbsp/2.5 gals)	1-2 qts of coarse spray per animal over whole body surface. For mange, scabies, ticks and lice, thoroughly wet animal. Repeat application in 10-14 days for mites
Horn flies, Face flies, Stable flies, Ear ticks	Low pressure sprayer	1 pt to 2-1/2 gals water (3 tbsp/2.5 gal)	1-2 qt spray per animal Spot treat back, face, legs and ears
Horn flies, Face flies, Stable flies	Backrubber/ Self oiler	1 pt to 10 gals diesel oil (3 tbsp/gal)	Keep rubbing device charged. Results improved by daily forced use.
Poultry			
Northern fowl mites, Lice	Sprayer	1 pt to 25 gals water (3 tbsp/2.5 gals)	1 gal of coarse spray per 100 birds per pen, particular attention to vent area.
Swine			
L. A. Mange	Sprayer	1 pt to 25 gals water (3 tbsp/2.5 gals)	Thoroughly wet animals including ears. For mange, repeat at 14 days.
For swine allow 5 days between last treatment and slaughter.			
Premises—in and around horse, beef, dairy, swine, sheep and poultry premises, animal hospital pens and barns and outside meat processing premises			
Horse flies, Stable flies, Little house flies (Fanni- ae) spp., Aids in control of ticks, mites, ants, spiders, mosquitoes, fleas, lice, etc.	Sprayer	1 pt to 10 gals water (3 tbsp/gal)	Spray to point of runoff or 1 gal per 700-1000 sq. ft.
	Overhead space spray system	1 pt to 10 gals diesel or mineral oil (3 tbsp/gal)	4 oz spray ; or 1000 cu. ft. of air space.
Do not apply dilutions for premises directly on livestock or poultry. Ensure that food and water are not contaminated by spray drift. For space spraying, do not remain in treated area and ventilate the area before reoccupying. Animals should be removed from area prior to treatment. Do not use in milk rooms. The use of any residual fly spray should be supplemented with proper manure management and general sanitation to reduce or eliminate fly breeding sites.			

Precautionary statements
Hazards to humans & domestic animals
Caution
Harmful if swallowed or absorbed through the skin.
Avoid contact with skin, eyes or clothing. Avoid
breathing spray mist.
Statement of practical treatment
If swallowed: Call a physician immediately DO NOT
INDUCE VOMITING unless under medical attention.
If on skin: Remove contaminated clothing and wash
skin thoroughly with soap and water.
If in eyes: Immediately flush eyes with plenty of
water. Get medical attention if irritation persists.
If inhaled: Remove victim to fresh air. Apply artificial
respiration if needed.
Note to physician: This product contains petroleum
distillates. Vomiting may cause aspiration pneumonia.
Environmental hazards
This product is extremely toxic to fish. Do not add
directly to water. Do not contaminate water by
cleaning of equipment or disposal of wastes.

Storage and disposal
Do not contaminate water, food or feed by storage or
disposal.
Storage: Store in cool, dry place.
Pesticide disposal: Wastes resulting from the use of
this product may be disposed of on site or at an
approved waste disposal facility.
Container disposal: Triple rinse (or equivalent). Then
offer for recycling or reconditioning, or puncture and
dispose of in a sanitary landfill, or by other procedures
approved by State and local authorities.

Statement of warranty
Solvay Animal Health, Inc. makes no warranty of
merchantability, fitness for any particular purpose, or
otherwise, expressed or implied concerning this
product or its use which extend beyond the use of
the product under normal conditions in accord with the
statements made on this label.
SOLVAY Animal Health, Inc.
Mendota Heights, MN 55120-1100 USA
Distributed from Charles City, IA 50616-0000 USA
4-000-0000
06 590000 1
Lot No.: 000

