

ACCE
 APR 7
 UNDER THE FEDERAL
 FUNGICIDE AND
 FOR ECONOMIC
 ED UNDER NO. 6
 TO APPL

CAREFULLY PRY OFF CAP. DO NOT USE FINGERS. RESEAL AT ONCE AFTER EACH USE AND BEFORE DISCARDING KEEP OUT OF THE REACH OF CHILDREN.

DANGER CAN CAUSE BURNS
 AVOID CONTACT WITH EYES, SKIN AND CLOTHES

Keep off wood, painted surfaces and floor coverings. Always use as is. Never make a solution. Do not use plunger. Keep hands and face away from drain after using this cleaner. Do not use with aluminum or galvanized utensils.

DIRECTIONS FOR USE

FOR SANITARY, FREE-FLOWING DRAINS: At least once a week pour into the drain 1/4 of this can of drain opener making certain there is at least one cupful of water in the drain. Follow with one cup of warm (not hot) water. If solution boils add cup of cold water. Let mixture work for 30 minutes then flush drain with warm water. Such regular use sanitizes drains by reducing odor-causing bacteria.

TO OPEN CLOGGED OR SLUGGISH HOUSEHOLD DRAINS: Remove all water possible from the inlet of drain. Remove strainer if possible. Slowly and carefully pour 1/3 the contents of this can into drain. Follow immediately with a cup of cold water. If solution boils, add another cup of cold water. Allow to remain for 1 hour then rinse with warm water. If drain is not completely clear, repeat treatment making certain that any standing water is removed before the second application. This water must not touch hands, skin or clothing. It should be dipped out with a china or glass cup (no aluminum) and poured down the nearest open drain. Wash cup immediately after using.

GENERAL USES

POWER DRAIN PIPE OPENER removes accumulations of lint, cloth, hair, slime, grease, vegetable matter and cigarette butts. Will dissolve stoppages in drains, bathtubs, kitchen sinks, dishwashing tubs, urinals, water closets, floor drains and grease traps. Will not injure pipe connections. Will not harm septic tanks.

Product will not clear stoppage caused by solid substances such as wood, iron, metal or broken glass. If stoppage is caused by any of these materials, be sure to advise plumber that **DRAIN PIPE OPENER** has been used so that he will not injure his hands or clothing.

ACTIVE INGREDIENTS:

Sodium Hydroxide	95.0%
Sodium Carbonate	1.0%
INERT INGREDIENTS	4.0%

POISON 

CONTAINS SODIUM HYDROXIDE

ANTIDOTES

EXTERNAL: Flood with water, then wash with vinegar.

INTERNAL: Give vinegar or juice of lemon, orange or grapefruit freely. Follow with olive oil, butter or other cooking oil.

EYES: Immediately flood with water for at least 15 minutes. Follow up with 5% Boric Acid Solution.

CALL PHYSICIAN AT ONCE

MADE IN U.S.A. • PRINTED IN U.S.A.
 U.S.D.A. REG. NO. 69-21

13 OUNCES NET WEIGHT
THE CHEMICAL PLUMBER



WARNING — KEEP OUT OF REACH OF CHILDREN. SEE BACK OF PANEL FOR ADDITIONAL CAUTIONS.

Packed for WY-TO CHEMICALS INC. CHICAGO, ILLINOIS

SIMAZINE 4% GRANULES

HERBICIDE

Granular Size 15/30

FOR WEED CONTROL IN ORCHARDS & NURSERY STOCK

ACTIVE INGREDIENT:

Simazine*(2-chloro-4, 6-bis (ethylamino)-s-triazine)-4%

INERT INGREDIENTS: - - - - -96%

GENERAL INFORMATION

Simazine 4G controls both broadleaf weeds and grasses. It is most effective as a pre-emergence herbicide and should be applied just prior to, or during, weed emergence. It is non-corrosive and non-inflammable. Simazine 4G acts through the roots and its effect on weeds is therefore dependent upon rainfall to move the chemical into the root zone. Excessively dry soil conditions and absence of rain may result in poor weed control.

DIRECTIONS FOR USE

Simazine 4G should be distributed uniformly over the soil surface by means of a Cyclone Spreader or other suitable mechanical application. For treatment of small areas, application by hand may be adequate.

NURSERIES: To control broadleaf weeds and grasses on Elm, Austrian Pine, Balsam Fir, Barberry, Blue Spruce, Box-elder, Bush Honeysuckle, Caragana, Cotoneaster, Douglas Fir, Honey Locust, Juniper, Multiflora Rose, Norway Spruce, Red Cedar, Red Oak, Red Pine, Rhododendron, Rose, Russian Olive, Scotch Pine, Siberian Elm, White Cedar, White Pine, Yew (Taxus spp.) Apply 50 to 75 lbs. of Simazine 4G per acre following and prior to weed emergence. Apply only to trees which have been established 1 year or more.

NON-BEARING APPLE, SOUR CHERRY, PEACH AND PEAR TREES FOR USE AROUND YOUNG TREES ONLY: For control of Annual Weeds and Grasses: Use 150 to 200 lbs. per acre. Apply early Spring (Feb. or March) during dormant season. Apply only to trees which have been established 1 year or more.

APPLES (bearing): To control Annual Weeds & Grasses: Apply 50 to 100 lbs. per acre in Spring before weeds emerge. Do not apply to foliage or fruit.

BLUEBERRIES, BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, AND RASPBERRIES: To control Broadleaf weeds and grasses: Apply 50 to 100 lbs. per acre in Spring before weeds emerge and before bud break, or broadcast in Fall. Do NOT apply when fruit is present. Do not apply to foliage.

CAUTION

Avoid contamination of irrigation ditches, sources of potable water and of beneficial plants. Do not apply to sandy soil.

Miller Chemical & Fertilizer Corp. makes no warranty of any kind, express or implied, concerning the use of this product. The use of this material being beyond our knowledge and control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, express or implied, as to the effects of such use, whether or not in accordance with directions or claimed so to be.

CAUTION: KEEP OUT OF REACH OF CHILDREN
NET WEIGHT 50 POUNDS

Manufactured By
MILLER CHEMICAL & FERTILIZER CORPORATION
Balto., Maryland 21215/Subsidiary of Alco Chemical Corp.

J.S. Patent 2,891,855
*Simazine is a trademark of Geigy Chemical Corp.

2/66 USDA Reg. No. 72-289 F.N. 2429 BATCH NO.

ACCEPTED