

70-11 PM-15 12/31/89 10/2

Use must be in accordance with
EPA Guidance for Phase 2
of the Insecticide Change in Status

distributed throughout the area. To reduce air movement as much as possible, it is preferable to close windows and doors before spraying; leave them closed for 10 minutes after spraying and then ventilate the area. Repeat daily, morning and afternoon, as needed. Do not apply as a barn spray if animals which have received direct application within 8 hours remain in the barn.

DO NOT CONTAMINATE FEED, DRINKING WATER, MILK AND MILKING EQUIPMENT.

CAUTION:

HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH THE SKIN. Do not breathe vapor or spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate water, feed or foodstuffs. Keep out of reach of children. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. Do not reuse container; destroy when empty. *S/et*

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

EPA Reg. No. 70-11
EPA Est. No. 70-KY-1

1025 N:01

RIGO

CONTAINS
PYRETHRINS
and
DDVP

**HORSE & STABLE
INSECTICIDE**
Ready-to-Use • Wipe on or Spray

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS

NET CONTENTS
1 LIQUID GALLON

Manufactured by—

B. → RIGO COMPANY, BUCKNER, KENTUCKY 40010

DIRECTIONS FOR USE

Rigo Horse and Stable Insecticide contains DDVP, piperonyl butoxide and pyrethrins. It is a ready-to-use spray for application directly to dairy cattle, beef cattle, hogs, sheep, goats, and horses to control horse flies, house flies, horn flies, stable flies, and mosquitoes, and to aid in control of face flies.

DIRECT SPRAYING OF LIVESTOCK—Apply as a finely atomized spray uniformly distributed over each animal at a rate of 1 to 2 fluid ounces per animal on dairy cattle, beef cattle, and horses, and at the rate of 1 fluid ounce per animal on hogs, sheep and goats. Reduce application rate by one-half for young animals. Use good spray equipment to spray each animal, preferably from the rear, directing the spray along both sides, including the belly, flanks, and withers, down each leg, over the back, and around the poll. Hold nozzle of sprayer 2 or 3 feet away from animal, so as to deposit droplets on the hair. **DO NOT WET THE HIDE** For milking dairy animals, apply the spray 20 to 30 minutes before beginning to milk, or after milking has been completed. **DO NOT SPRAY UDDERS OF LACTATING DAIRY ANIMALS. DO NOT APPLY IN EXCESS OF RATES STATED PER ANIMAL.** Repeat daily as needed.

BARN SPRAYING—To control house flies in dairy barns, milk rooms, loading sheds, livestock barns, stables, and poultry houses, apply 2 fluid ounces per 1,000 cubic feet as a finely atomized spray, uniformly

ATTACHMENT 1

A. ACTIVE INGREDIENTS:

Dichlorvos: 2,2-dichlorovinyl dimethyl phosphate	0.18%
Related compounds	0.02%
Pyrethrins	0.03%
Technical piperonyl butoxide *	0.30%
Inert ingredients:	99.47%
Total	<u>100.00%</u>

* Equivalent to 0.24% (butylcarbityl) (6-propylpiperonyl) ether and 0.06% related compounds.

B. Rigo Company
A Division of Wilbur-Ellis Company
Buckner, Kentucky 40010