

SILVERFISH and FIRE BRATS: Thoroughly treat areas behind shoeing book cases and storage areas and any other infested cracks or hiding places. Also apply locally on exposed surfaces where Silverfish crawl when they come out of hiding. Use 6 tablespoons per gallon of water. For spot application only.

CARPET BEETLES: Apply to infested areas on the Floor, Baseboards and Closet Shelves and work material into cracks where insects may be found. Treat underside of carpets. Use 6 tablespoons per gallon of water. For spot application only.

WASPS: Apply a liberal application of the nesting sites of Wasps, such as underneath eaves, porches, sheds, and garages. Apply after dark when insects have gone to rest, so as to avoid possible stings. Use 6 tablespoons per gallon of water.

BOX ELDER and HOUSE CRICKETS: Apply to outside areas where they crawl or gather in masses. Treat thoroughly entrances and in closets around baseboards and other places where they may hide. Use 6 tablespoons per gallon of water. For spot application only.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

**ENVIRONMENTAL CAUTIONS:**

This product is toxic to fish, birds and other wildlife. Birds and other wildlife treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**WARNING:** May be fatal if swallowed! Contact with skin can cause toxic symptoms. Avoid breathing Spray Mist. Keep out of reach of children. Read directions carefully and note warnings before using. In case of skin contact wash with soap and water. Avoid contamination of feed and feedstuffs. Food utensils such as tablespoons or measuring cups should not be used for food purposes after use with pesticides.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Keep children and pets off treated areas until this material is washed into soil and grass is dry.

Do not reuse empty container. Destroy if leaking, perforating or cracking. Bury or discard in a safe place.

2081-725



74%  
CHLORDANE  
EMULSIFIABLE

ACTIVE INGREDIENTS:

Technical Chlordane 74  
Petroleum Distillate 6

EPA Reg. No. 75-381  
EPA Est. No. 254-342-01

INERT INGREDIENTS:

\*Equivalent to 49-4-Octa chloro-cyclohexane, 2,3,4,5-tetra-  
hydro-dane and 29-6-Pentachlorocyclohexene.

NET CONTENTS  
ONE LIQUID QUART

KEEP OUT OF REACH OF CHILDREN.

SEE SIDE PANEL FOR DIRECTIONS

KILLS

For use in home  
yards, garbage  
cans, dumps.

**DIRECTIONS**

ROACHES or WATERBUGS, SPIDERS and ANTS INSIDE BUILDINGS: Thoroughly treat at rate of 3 tablespoons per gallon of water infested cracks, baseboards, under sinks and drainboards, door sills and along runways. Use as spot treatment only.

NOTE: Keep away from food surfaces or containers. Avoid food contamination.

ARMY WORMS, SOD WEBWORMS, JAPANESE BEETLE GRUBS, CHINCH BUGS: Spray thoroughly 4 teaspoons in 1 gallon of water per 150 square foot area for Lawns, Nursery Stocks and Ornamental Plants. (Not effective on Chinch Bugs in Florida.)

CUTWORMS, [Brachythrus] on Camellias, Azaleas, Rhododendrons, Primrose and many other ornamentals: Spray generously at 4 teaspoons per gallon of water around base of plants working into top 4 inches of soil.

**TERMITE CONTROL:**

Prepare a 1% emulsion by mixing 3 tablespoons per gallon of water and use mixture as follows:

SLAB-TYPE CONSTRUCTION: (a) Apply 1 gallon per 10 sq. ft. as an over-all pre-construction treatment under the slab and under attached porches. (b) Apply 2 gallons per 5 linear feet to crucial areas under the slab as pre-construction treatment, such as along the inside of foundation walls and around utility entrances and interior partition foundation walls. (c) Apply along the outside of the foundation as indicated below.

BUILDINGS HAVING BASEMENTS OR CRAWL SPACES: Dig a trench along the outside of the foundation walls. In brick or hollow block or concrete foundation, dig a trench to, but not below, the footing. Then as the trench is refilled, treat the soil at the rate of 4 gallons per 10 linear feet for each foot of depth. A trench 3 feet deep would require 12 gallons per 10 linear feet.


The objective is to establish a treated soil barrier which will prevent termite entry. Other critical areas may also require treatment. It is suggested that you consult the State Extension Service for advice.

WOOD TREATMENT: There may be occasion when it is desirable to protect wood from termites, such as fence posts or mail box posts. Follow the directions to make a 1% Termi-Ded emulsion, however, oil (such as used fuel oil) may be used instead of water as a diluent. Soak a portion of the post which will be in the ground for 23 hours. Wear protective clothing and rubber gloves when handling the freshly treated lumber.

RIGO COMPANY



BUCKNER, KENTUCKY 40010



Comment

ACCEPTED WITH COMMENTS