

**DIRECTIONS  
CONTINUED**

**STRAWBERRY (4):** Meadow Spittlebug, Strawberry Aphid—4 ozs. per 500 sq. ft. Do not apply at intervals of less than 15 days or more than twice during any 35 day period when fruit is present.

**SUGAR BEETS:** Green Peach Aphid, Sugar Beet Webworm—4 ozs. per 500 sq. ft. Meadow Spittlebug—2 ozs. per 500 sq. ft. Do not feed sugar beet tops to livestock.

**TOBACCO (5)** Cabbage Looper, Green Peach Aphid, Tobacco Budworm, Tobacco Flea Beetle, Tobacco Hornworm—SEED BED 1½ lbs. per 100 sq. yds. FIELD: 15 to 24¼ lbs. per acre. Cutworms, Pillbugs, Slugs (FLORIDA-GEORGIA SHADE GROWN TOBACCO AREA ONLY) .375 to .56 lbs. per 100 sq. yds. plant bed. Direct dust at hiding places of these pests.

**TOMATOES (1):** Blister Beetles, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid—2 ozs. per 500 sq. ft. Cabbage Looper—3 ozs. per 500 sq. ft. Stink Bugs—4 ozs. per 500 sq. ft. Tomato Fruitworm, Yellow-striped Armyworm—4 ozs. per 500 sq. ft. (For both field and greenhouse use). Tomato Hornworm—4 ozs. per 500 sq. ft. Tomato Russet Mite—4 ozs. per 500 sq. ft. Whitefly—2 ozs. per 500 sq. ft. (For both field and greenhouse use).

**CAUTION: KEEP OUT OF REACH OF CHILDREN.** Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Workers entering treated areas within 24 hours of application should wear protective clothing.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Keep out of lakes ponds and streams. Do not apply to any area not specified on the label.

Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

**KILL-KO**

**KILL-KO**

**KILL-KO**

**KILL-KO**

**NET WT. 4 LBS.**

**KILL-KO®**

**THIODAN®  
4 DUST**

**FOR USE ON TOBACCO**

**ACTIVE INGREDIENT:**

\*Endosulfan (Hexachlorohexahydromethano-2,4,3-bensodioxathiepin oxide) ... 4.00%

**INERT INGREDIENT:** ..... 96.00%  
100.00%

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**  
See back panel for additional cautions.

\*Thiodan U. S. PATENT NO. 2,799,685  
Thiodan is a registered trademark of Canadian Hoechst Limited.

MANUFACTURED BY  
**RIGO Chemical Company**  
NASHVILLE, TENN. 37202

Acceptance stamp on pg. 3 of 3

70-126

11-2-71

1043

## DIRECTIONS CONTINUED

**LETTUCE, HEAD (14):** Cabbage Looper, Diamond Back Moth Larvae, Green Peach Aphid, Imported Cabbageworm—4 ozs. per 500 sq. ft. Do not make more than 3 applications after thinning. Remove wrapper leaves at harvest. Do not feed crop refuse to livestock.

**MELONS (0):** Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle—4 ozs. per 500 sq. ft. Cabbage Looper, Omnivorous Leaf Roller—6 ozs. per 500 sq. ft. Rindworm (on Water melon only)—4 ozs. per 500 sq. ft.

**ORNAMENTALS (Greenhouse and Out-of-doors):** Aphids, Whitefly—Dust lightly but evenly. On Chrysanthemums best results will be obtained if applied before plants flower. Do not use on Birch trees. In Greenhouse, wear mask or respirator approved by USDA for pesticide protection.

**PEAS (SEED PEAS ONLY):** Pea Aphid, Pea Weevil—4 ozs. per 500 sq. ft. Do not feed treated vines or threshings to livestock.

**PEPPERS (1):** Flea Beetle, Green Peach Aphid, Hornworms—4 ozs. per 500 sq. ft. Pepper Maggot—2 ozs. per 500 sq. ft.

**POTATOES (0):** Aphids, Armyworms, Colorado Potato Beetle, Green Stink Bug, Leaf-footed Bug, Plant Bugs, Potato Flea Beetle, Potato Leafhopper, Potato Tuberworm—4 ozs. per 500 sq. ft. European Corn Borer, False Chinch Bug, Potato Psyllid—4 ozs. per 500 sq. ft. Three-lined Potato Beetle—2 ozs. per 500 sq. ft. Whitefly—3 ozs. per 500 sq. ft.

**PUMPKINS (0):** Aphids, Cucumber Beetles, Squash Vine Borer—4 ozs. per 500 sq. ft. For Vine Borer apply weekly to flower buds, stems and vines beginning when moths first appear.

**SAFFLOWER:** Green Peach Aphid—4 ozs. per 500 sq. ft. Do not apply after flower heads open.

**SOYBEANS:** Mexican Bean Beetle, Stink Bug—2½ ozs. per 500 sq. ft. Do not apply after pods begin to form. Do not feed treated threshings to livestock.

CONTINUED ON OTHER SIDE PANEL

## DIRECTIONS FOR USE

Apply when insects first appear and repeat as required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in ( ) following crop.

**APPLES (30):** Apple Aphid—6 ozs. per 500 sq. ft. Unlimited number of applications. Do not feed pomace to livestock.

**APPLES (21):** Apple Aphid—6 ozs. per 500 sq. ft. Do not make more than 3 applications during the fruiting period. Do not feed pomace to livestock.

**APRICOTS (30), NECTARINES (30), PEACHES (30):** Green Peach Aphid—6 ozs. per 500 sq. ft. Do not make more than 2 applications during the fruiting period.

**ARTICHOKES (7):** Artichoke Aphid, Artichoke Plume Moth, Green Peach Aphid—4 ozs. per 500 sq. ft.

**BEANS (DRY BEANS ONLY) (3):** Bean Aphid, Bean Leaf Skeletonizer, Mexican Bean Beetle—4 ozs. per 500 sq. ft. Cucumber Beetle—3 ozs. per 500 sq. ft. Do not apply more than 3 times per season. Do not apply within three days of harvest. Do not feed treated threshings to livestock or allow to graze in treated fields. Do not use on Lima Beans that are to be removed from the field for processing. Do not feed to livestock.

**BEANS (SUCCULENT):** Cucumber Beetle—3 ozs. per 500 sq. ft. Mexican Bean Beetle—3½ ozs. per 500 sq. ft. Do not apply after pods begin to form. Do not apply more than three times per season. Do not apply within three days of harvest. Do not feed treated threshings to livestock or allow to graze in treated fields. Do not use on Lima Beans that are to be removed from the field for processing. Do not feed to livestock.

**BROCCOLI (7) CABBAGE (7):** Cabbage Aphid, Cabbage Looper, Cross-Striped Cabbageworm, Diamond Back Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbage Worm, Stink Bug—4 ozs. per 500 sq. ft.

**CAULIFLOWER:** Cabbage Looper, Cross-Striped Cabbageworm, Diamond Back Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbageworm, Stink Bug—4 ozs. per 500 sq. ft. Do not apply after edible parts begin to form. Close trim outer leaves prior to marketing.

**COTTON:** Boll Weevil—4½ ozs. per 500 sq. ft. Bollworm, Cabbage looper, Lygus Bugs, Stink Bugs—4 ozs. per 500 sq. ft. Cotton Leaf Performator—4 ozs. per 500 sq. ft. Thrips—6 ozs. per 500 sq. ft. Do not apply after bolls open. Do not feed treated plants to meat or dairy animals.

**CUCUMBERS (0), SQUASH (SUMMER AND WINTER) (0):** Aphids, Cucumber Beetles, Squash Vine Borer, Striped Cabbage Flea Beetle—4 ozs. per 500 sq. ft. For Squash Vine Borer apply weekly to flower buds, stems and vines beginning when moths first appear.

**EGGPLANTS (1):** Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, Green Stink Bug, Whitefly—2¼ ozs. per 500 sq. ft.

**FILBERTS (0):** Filbert Aphid, Filbert Leaf Roller—6 ozs. per 500 sq. ft.

**GRAPES (7):** Grape Leafhopper—6 ozs. per 500 sq. ft. Do not apply to Concord variety as severe injury is likely to occur.

**IRIS:** Iris Boere—Dust soil surface of beds thoroughly in spring when growth starts. Three applications are needed at 2-week intervals.

USDA REG. NO. 70-126

CONTINUED IN BOTH SIDE PANELS

2013

**ACCEPTED**  
**NOV 2 1971**  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. .... SUBJECT  
TO ATTACHED COMMENTS.

70-126

Faint, illegible text, likely bleed-through from the reverse side of the page.