

Provide one face oiler for each 50 head of cattle. Provide one back rubber for each 50 head of cattle. For effective results the face rubber and back rubber must be accessible continuously and must be refilled as necessary.

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid inhalation of mist and contact with skin. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention.

Wash thoroughly with soap and water after handling and before eating or smoking.

Do not contaminate feed, water or foodstuffs. Keep out of reach of children.

If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.

Combustible liquid, do not use, spill, pour or store near heat or open flames.

Rinse empty container thoroughly with water and discard it.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

1151-734

EPA Reg. No. 70-130

EPA Est. No. 70-KY-1

RIGO COMPANY

Buckner, Kentucky 40010

70-130



**FLY SPRAY
FOR DAIRY AND
BEEF CATTLE**

CONTROLS:

- HORN FLIES
- FACE FLIES
- STABLE FLIES
- HOUSE FLIES
- MOSQUITOES
- LICE

**CONTAINS 1.0% CIODRIN® INSECTICIDE
AND 0.25% VAPONA® INSECTICIDE**

ACTIVE INGREDIENTS

Petroleum Hydrocarbons 38.57%

Dimethyl phosphate of alpha-methylbenzyl 3-hydroxy-cis-crotonate* 1.00%

2,2-dichlorovinyl dimethyl phosphate 0.23%**

Related Compounds 0.02%**

INERT INGREDIENTS 0.18%

TOTAL 100.00%

*Ciodrin® Insecticide

**Equivalent to 0.25% Vapona® Insecticide

CIODRIN® AND VAPONA® ARE REGISTERED
SHELL TRADEMARKS

**NET CONTENTS
ONE LIQUID GALLON**

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panel for additional cautions

DIRECTIONS

FOR USE AS A CATTLE SPRAY

Ready to use—Do not dilute

For control of horn fly, face fly, stable fly, house fly and mosquito on cattle, apply a light application daily as a fine mist. Spray to thoroughly cover all parts of the animal, including the legs. Do not apply in excess of 2 fluid ounces per animal per day.

Do not wet the skin due to possible skin irritation. Do not apply regularly to calves under six months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

**FOR CONTROL OF LICE ON
DAIRY AND BEEF ANIMALS**

Make 1-2 applications of 3-3½ ounces of Kill-Ko FLY SPRAY at 14 day intervals as a fine mist spray to thoroughly cover all parts of the animal. If material is applied by fogging or misting equipment, the maximum spraying time per animal is 20 seconds.

**FOR USE IN FACE RUBBERS
& BACK RUBBERS**

Dilute before using

For the control of horn flies and face flies on dairy and beef cattle.

When used in face rubber and back rubber devices this product must be diluted three quarts of this insecticide with one quart of No. 2 fuel oil or diesel oil to make one gallon of face and back rubber solution.