

EPA Est. No. 70-KY-1

R ->

RIGO COMPANY

Buckner, Kentucky 40010

PRECAUTIONARY STATEMENTS

2 5 4

۶.

·: .

Hazards to Humans and Domestic Animals

2. ,-- '

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid inhalation of mist and contact with skin. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention.

Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water or foodstuffs.

Neep out fo reach of children.

If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.

Environmental Hazards

This pesticide is toxic to fish, birds, and aquatic invertebrates. Do not apply directly to water or wetlands (swamps, marshes, bogs, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Combustible liquid, do not use, spill, pour or store near heat or open flames.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

"Wear chemical resistant gloves, long-sleeved shirt, long-legged pants, shoes and socks when handling the product. Do not allow children or pats to enter treated rooms until thoroughly ventilated, or to contact treated surfaces until sprays have dried."

FOR USE AS A CATTLE SPRAY

Ready to use-Do not dilute

For control of horn fly, face fly, stable fly, house fly and mosquito on cattle, apply a light application daily as a fine mist. Spray to thoroughly cover all parts of the animal, including the legs. Do not apply in excess of 2 fluid ounces per animal per day.

Do not wet the skin due to possible skin irritation. Do not apply regularly to calves under six months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

FOR CONTROL OF LICE ON Dairy and beef animals

Make 1-2 applications of 3-3½ ounces of Kill-Ko FLY SPRAY at 14 day intervals as a fine mis spray to thoroughly cover all parts of the animal If material is applied by fogging or misting equipment, the maximum spraying time per animal is 20 seconds.

FOR USE IN FACE RUBBERS & BACK RUBBERS

Dilute before using

6/88

For the control of horn flies and face flies on dairy and beef cattle.

When used in face rubber and back rubber devices this product must be diluted three quarts of this insecticide with one quart of No. 2 fuel oil or diesel oil to make one gallon of face and back rubber solution.

Provide one face oiler for each 50 head of cattle. Provide one back rubber for each 50 head of cattle. For effective results the face rubber and back rubber must be accessible continuously and must be refilled as necessary.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container only, in a cool, dry location inaccessible by children or domestic animals. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Triple rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved state and local procedures.

use and/or handling Buyer assumes all NUTICE: Seller makes no warranty, expressed or implied, concerning the when such the label. material use of this product other than indicated on of this to label instructions and/or handling risk of use contrary 8

3 7

4

ATTACHMENT 1

A. ACTIVE INGREDIENTS:

• •

Dimethyl phosphate of alpha-methylbenzyl		
3-hydroxy-cis-crotonate *	1.00%	
Dichlorvos [2,2-dichlorovinyl dimethyl phosphate]	0.23% *	:*
Related Compounds	0.02% *	*
Inert Ingredients:	98.75%	
-		

Total 100.00%

4 + 4

* Ciodrin Insecticide ** Equivalent to 0.25% Vapona Insecticide

B. Rigo Company A Division of Wilbur-Ellis Company Buckner, Kentucky 40010