



LONG TERM CONTROL:

Bait continuously using KILL-KO THRO PAC. Stale, damp or moldy bait should be replaced with fresh bait promptly. Treated bait should be placed at locations not accessible to children, pets, wildlife and domestic animals or in tamper-proof bait boxes.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Keep away from children, pets, wildlife and domestic animals. In case baits are eaten by humans, call physician at once. Treat as for overdose of Dicoumarol®. **3051-7111**


MAR 5 1973
INDEXED
SERIALIZED
FILED
FBI - MEMPHIS
73
SUBJECT TO APPLICABLE ORDINANCES.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See back panel for additional cautions

ACTIVE INGREDIENTS: Diphacinone 2-diphenylacetyl — 1, 3-Indandione .005%
INERT INGREDIENT: 99.995% NET WEIGHT 4 OZ.

MANUFACTURED BY
RIGO CHEMICAL COMPANY, NASHVILLE, TENN. 37202
EPA REG. NO. 70-139



KILL-KO, THRO PAC Rat Killer

DIRECTIONS:

RATS:

Place 1 to 2 KILL-KO THRO PAC next to burrows in runways or where rats are accustomed to feeding. Check these treated bait stations daily to assure an uninterrupted supply of bait for a period of not less than 12 days. Continue baiting until all signs of feeding have stopped. One KILL-KO THRO PAC is effective for one or two rats.

MICE:

Place 1 ounce of KILL-KO THRO PAC in a shallow dry container next to burrows in runways or where mice are accustomed to feeding. It is recommended that KILL-KO THRO PAC be placed at intervals of 8 to 12 feet. Check treated bait stations daily to assure uninterrupted supply of KILL-KO THRO PAC for a period of not less than 15 days.