

RIGO'S BEST

THIODAN

INSECTICIDE

WARNING

Hazardous if swallowed, inhaled or absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water, for eyes get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking, wear clean clothing. During commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U.S. Bureau of Mines for Endosulfan protection. Do not allow to drift to areas occupied by unprotected humans or beneficial animals. Do not use for aircraft application.

Workers entering treated areas within 24 hours of applications should wear protective clothing.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding in treated areas will be killed at rates listed.

Do not apply when weather conditions

are not from areas listed. Do not apply where runoff is likely to occur. This product is toxic to bees and should not be applied when bees are actively visiting the area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Store in original container in a cool, dry location away from heat, flame, direct or indirect sunlight. Do not store at temperature below 20 F. Do not reuse empty container. Wrap container and put in trash collection.

Apply this product only as specified on this label.



NET CONTENTS 1 QUART

MANUFACTURED BY RIGO COMPANY,
BUCKNER, KENTUCKY 40010

B



A }

ACTIVE INGREDIENTS
Endosulfan (Hexachlorocyclohexane)
hydromethane-2,4
3-benzodioxathepoxide
24.0% Heavy Aromatic
Naphtha 69.0% INERT
INGREDIENTS 7.0%

Thiodan is a registered trademark of Canadian Hoechst Limited
EPA Reg. No. 70-142 EPA Est. No. 70-KY-1

WARNING: KEEP OUT OF REACH OF CHILDREN

← SEE BACK PANEL FOR ADDITIONAL CAUTIONS

BEST AVAILABLE

A. ACTIVE INGREDIENTS:
Endosulfan (Hexachlorohexahydromethano-2,4,3-
benzodioxathiepin oxide.....24.0%

INERT INGREDIENTS*:. 76.0%

TOTAL 100.0%

*Contains xylene range aromatic solvent.

B. Rigo Company
A Division of Wilbur-Ellis Company
Buckner, KY 40010

Attachment (70-142)

- A. ACTIVE INGREDIENTS:
Endosulfan (Hexachlorohexahydromethano-2,4,3-
benzodioxathiepin oxide.....24.0%
- INERT INGREDIENTS*:. 76.0%
- TOTAL 100.0%

*Contains Heavy Aromatic Naphtha.

- B. Rigo Company
A Division of Wilbur-Ellis Company
Buckner, KY 40010

Looper - FIELD - 2 to 4 tbs. SEED BED - Use 2 tbs. in 3 gal. of water and apply 3 gal. per 50 sq. yd. Green June Bug Larvae - Use 1 tbs. in 3 gal. of water and drench plant bed with 1 gal. spray per sq. yd. Slink Bugs - Use field application of 4 to 6 tbs. in 3 gal. of water per 1375 sq. ft.

TOMATOES (Field and Greenhouse) (1): Slater Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid - 2 tbs. in 3 gal. of water. Hornworms - 2 to 4 tbs. in 3 gal. of water. Tomato Russet Mite, Tomato Fruitworm, Yellow-Striped Armyworm - 4 tbs. in 3 gal. of water. Slink Bugs - 3 to 4 tbs. in 3 gal. of water. Cabbage Looper - 3 tbs. in 3 gal. of water. Whitefly - 2 tbs. in 3 gal. of water.

ORNAMENTALS (Greenhouse & out of doors): Aphids, Cyclamen Mite, Whitefly, Rose chafar - 2 tbs. in 3 gal. of water. On Chrysanthemums, best results will be obtained if applied before plants flower. Do not apply to "Bonnaton Deluxe", "Fred Showman" and "White Knight" chrysanthemums as injury may result. Black Vine Weevil - 4 tbs. in 3 gal. of water. Spray thoroughly and drench soil under plants. Apply when beetles first appear. Repeat in 14 days. Do not use on birch trees. In greenhouse applications wear mask or respirator passed by USDA for Endosulfan protection.

SHADE TREES (Except Birch): Aphids - 2 tbs. in 3 gal. of water.

ELM: Aphids - 2 tbs. in 3 gal. of water. Elm Leaf Beetle (New England area only) 4 tbs. in 3 gal. of water. Apply when 50% of the overwintering eggs have hatched.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

RIGO COMPANY
BUCKNER, KENTUCKY 40010

4-68

RIGO'S BEST
THIODAN
INSECTICIDE CONCENTRATE

Controls over 50 major
crop & garden pests
including Colorado
Potato Beetle.

SEE CONTAINER LABEL FOR
INGREDIENTS STATEMENT AND
COMPLETE PRECAUTIONARY
STATEMENTS.

Directions for Use

It is a violation of Federal law to use this product
in a manner inconsistent with its labeling.

DIRECTIONS

Apply the listed amount per 1375 sq. ft. when insects first appear and repeat as required to maintain effective control. Use in sufficient water for thorough coverage unless otherwise noted for listed crops. Coverage of upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in () following the crop.

BEANS: Succulent and dry, except LIMAS (3) Black Bean Aphid, Bean Leaf Skeletonizer, Cucumber Beetles, Green Slink Bug, Mexican Bean Beetle - 2 to 4 tbs. in 3 gal. water per 1375 sq. ft. Do not exceed 3 applications when bean pods are present. Do not feed freshings or allow livestock to graze on treated fields. Do not exceed 3 applications per season.

BROCCOLI (7), CABBAGE (7), BRUSSELS SPROUTS (14), CAULIFLOWER (14): Cabbage Aphid, Cabbage Looper, Cross-striped Cabbage Worm, Diamondback Moth Larvae, Flea Beetle, Harlequin Bug, Imported Cabbage Worm, Slink Bug - 3 to 4 tbs. per 3 gal. water per 1375 sq. ft.

CUCUMBERS (6), MELONS (0), SUMMER & WINTER SQUASH (0)

Aphids, Cucumber Beetles, Striped Cabbage Beetle, Melonworm, Pickleworm, Rindworm (on watermelons), Squash Beetle, Squash Bug, Squash Vine Borer - 2 to 4 tbs. per 3 gal. water per 1375 sq. ft. For Vine Borer control, apply weekly to flower buds, stems and vines beginning when moths first appear. For Cabbage Looper and Omnivorous Leaf Roller on Melons, use 4 tbs. per 1375 sq. ft. in 3 gallons of water.

LETTUCE HEAD (14): Cabbage Looper, Green Leaf Aphid, Diamondback Moth Larvae, Imported Cabbage Worm - 3 to 4 tbs. per 3 gal. water per 1375 sq. ft. Do not make more than 3 applications after thinning. Remove wrapper leaves at harvest. Do not feed crop refuse to livestock.

PEPPERS (4): Green Peacan Aphid, Flea Beetle, Hornworms, Pepper Maggot - 2 to 4 tbs. per 1375 sq. ft. in 3 gal. of water.

POTATOES (6): Aphids, Colorado Potato Beetle, Plant Bugs (including Shield-shaped Bugs), Potato Flea Beetle, Potato Leaf Hopper, Potato Tuberworm, Southern Armyworm, Whitefly - 2 to 4 tbs. per 1375 sq. ft. in 3 gal. of water. European Corn Borer, Potato Psyllid - 3 to 4 tbs. per 1375 sq. ft. in 3 gal. of water. False Chinch Bug - 4 tbs. per 1375 sq. ft. in 3 gal. of water.

SPELNACH (21): Crown Mite, Green Peach Aphid - 3 tbs. per 1375 sq. ft. in 3 gal. of water. Do not exceed one application per season.

STRAWBERRIES (4): Cyclamen Mite - 2 tbs. per 1375 sq. ft. in 3 gal. of water. Do not apply at intervals of less than 35 days when fruit is present. Meadow Spittlebug, Strawberry Aphid, Tarnished Plant Bug - 2 tbs. per 1375 sq. ft. in 3 gal. of water. Do not apply at intervals of less than 15 days or more than twice during any 35 day period when fruit is present.

SWEET CORN (Fresh Vegetables use only) (6): Corn Leaf Aphid - 4 tbs. in 3 gal. of water per 1375 sq. ft. Corn Earworm - 6 tbs. in 3 gal. of water per 1375 sq. ft. Apply a 2 to 3 day intervals beginning when silks first appear and continuing until they begin to dry. Do not exceed 5 applications. Do not apply to sweet corn to be processed. Do not feed treated forage or ensilage to livestock or allow to graze in treated fields.

TOBACCO (5) Flea Beetle, Hornworm, Budworm, Green peach Aphid, Cabbage

BEST AVAILABLE COPY