General Precautions:

Do not spray delicate new growth or sensitive plants such as cyclamen, lantana, maiden hair fern or dull-leafed varieties of roses. Do not use on tomatoes, vegetables or other food plants. Do not spray on plastics or asphalt tile.


CAUTION: KEEP OUT OF REACH OF CHILDREN

Contents harmful if swallowed. The spray from this product may be a skin irritant and sensitizer. Avoid skin contact. In case of skin contact, wash with soap and water. Avoid breathing spray mist. Cover or remove exposed food, utensils and serving areas before spraying. Cover or remove fish for one hour.

WARNING: FLAMMABLE: Do not spray near fire or open flame. CONTENTS UNDER PRESSURE: Store upright in a cool place (not over 120°F). Do not place on hot surface. Do not puncture can. Never throw in fire or incinerator.

Manufactured for

RIGO CHEMICAL COMPANY Nashville, Tenn 37202




DIRECTIONS FOR USE

Remove plastic cap. Hold container upright. Point opening away from you and press button. Use indoors or outdoors.

For aphids, mites, thrips (exposed), white fly, lace bugs, climbing cutworms and Japanese beetles on ornamental plants and roses: more dispenser about 18" from foliage. Spray plant foliage lightly but thoroughly to wet both sides of leaves and insects that are present. Repeat treatments as necessary. Do not overspray as burning may result.

For black spot and powdery mildew on roses: Hold dispenser about 18" from foliage. Spray plant foliage lightly but thoroughly to wet both sides of leaves. Apply when new growth appears in the spring and repeat at 7 to 10 day intervals throughout the season. Do not overspray, as burning will result.

For use indoors apply sparingly in short bursts. For best results outdoors, spray in evening or on cloudy day when air is still. Use slow sweeping motion.





Kills aphids, mites, thrips (exposed), white fly, lace bugs, climbing cutworms and Japanese beetles on ornamental plants and roses. Controls black spot and powdery mildew on roses.

ACTIVE INGREDIENTS:

ary reitorins		.020%
this tell an cloheptene disc	arbor milde	300°°
Register Tonic		.100 ' -
ther opinites as		.200° -
to the track, how therein ear		.300* ~
Lope Podable Can	Chitting and the	120°
The try borry the control	oheny crutorate	1130
rich that then and the	vatives, chiefy	
" methy hopty" 4 mid sidrog !	henril	012°n
war early date ate		11.5%
INERT INGREDIENTS:		98.72
	7	7 ()

Tota1 (1961)

The second control of the second

The first of the section of the section of the contract of the and the second of the second of the second

AUTION FEED OUT OF REACH OF CHILDREN

The second of th

A CONTRACTOR OF THE STATE OF TH