

4-26 74

70-155

**KILL-KO**

# FLY SPRAY CONCENTRATE

EMULSIFIABLE CONCENTRATE  
DAIRY AND LIVESTOCK SPRAY

Contains CIODRIN® and VAPONA® Insecticides

ACTIVE INGREDIENTS	BY WEIGHT
Dimethyl phosphate of Alpha-methylbenzyl 3-hydroxy-cis-crotonate	10.0%
2, 2-dichlorovinyl dimethyl phosphate	2.3%
Related Compounds	0.2%
Heavy Aromatic Naphtha	77.0%
<b>INERT INGREDIENTS</b>	<b>10.5%</b>

\*Equivalent 2.5% Vapona® Insecticide  
CIODRIN® and VAPONA®  
® are Registered SHELL Trademarks

**NET CONTENTS: 1 FLUID GALLON**

**WARNING: Keep Out of Reach of Children**

See panel for additional caution.

## DIRECTIONS FOR LIVESTOCK PEST CONTROL

For Control of House Flies, Face Flies, and Stable Flies In and Around Livestock Buildings.

Mix 1 pint of KILL-KO FLY SPRAY CONCENTRATE per 1 1/2 gallons of water (1% concentration). Spray premise such as inside and outside barn walls (including dairy barns), eaves, fences and other resting or breeding places at least twice monthly using approximately 1 gallon of finished spray per 1000 square feet (40' x 25'). Improved house fly and face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6-10 gallons of spray.

## PRECAUTIONS IN USING

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS  
WARNING: HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. DO NOT BREATHE SPRAY MIST. KEEP AWAY FROM HEAT AND OPEN FLAME.

Wash thoroughly with copious water after handling and before eating and smoking. Wear clean clothing. In case of spilling on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water. If eyes get medical attention. Atropine is antitoxin. 2-PAM is an antidote and may be administered in conjunction with atropine.

U.S. DEPARTMENT OF AGRICULTURE  
BUREAU OF ENTOMOLOGY  
WASHINGTON, D.C. 20250  
KILL-KO FLY SPRAY CONCENTRATE  
REGISTERED TRADEMARK SHELL OIL CORPORATION  
© 1974 SHELL OIL CORPORATION



# FLY SPRAY CONCENTRATE

## EMULSIFIABLE CONCENTRATE DAIRY AND LIVESTOCK SPRAY

Contains Chlorfenvinphos and DAPONA® Insecticides

BY WEIGHT

Chlorfenvinphos	10.0%
DAPONA® (Dimethyl phosphite)	2.3%*
Other ingredients	87.7%
<b>Total</b>	<b>100%</b>

NET CONTENTS: 1 PLOID GALLON

### DIRECTIONS FOR LIVESTOCK USE

For Control of House Flies, Face Flies, and  
Around Livestock Buildings.  
Mix 1 pint of KILL-KO FLY SPRAY CONCENTRATE  
of water (1% concentration). Spray premise  
outside barn walls (including dairy barns), eaves  
resting or breeding places at least twice monthly.  
mately 1 gallon of finished spray per 1000 square  
improved house fly and face fly control may be  
achieved.

### PRECAUTIONS IN USE

KEEP OUT OF REACH OF CHILDREN AND DO  
NOT SWALLOW. IRRITANT IF SWALLOWED. INHALATION  
IRRITANT. SKIN IRRITANT. GET IT OFF IN EYES,  
IMMEDIATELY FLUSH WITH WATER. AVOID CONTACT WITH  
MILK AND MILK PRODUCTS.

Apply this spray to the exterior walls of buildings  
and to the eaves and roof of buildings. Do not spray  
directly on animals. Do not spray in the presence of  
livestock. Do not spray in the presence of children.  
Do not spray in the presence of milk or milk products.  
Do not spray in the presence of food or feed.