

DIRECTIONS FOR LIVESTOCK PEST CONTROL

For Control of Stable Flies, Horn Flies, House Flies, Lice and Lone Star Ticks on Dairy Cattle, Beef Cattle, Sheep, Goats and Hogs.

Mix 1 pint of KILL-KO FLY SPRAY CONCENTRATE per 1 1/2 gallons of water. Spray to thoroughly cover all parts of the animals, including the legs, using 1-2 pints of spray per large animal and proportionately less for smaller animals.

For pen or corral operations involving spray application to livestock with high pressure equipment, do not exceed a 0.5% concentration. Mix 1 pint of KILL-KO FLY SPRAY CONCENTRATE per 3 gallons of water (0.5% concentration). Spray to thoroughly cover the animal's body and legs (avoid excessive spraying of the head) using 1-2 quarts of spray per large animal and proportionately less for small animals. Repeat above application as necessary to maintain control, but not more often than once every 7 days. Improved face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6-10 gallons of spray. Repeat above application as necessary to maintain control, but not more often than once every 7 days.

Low Volume Application for Control of Face Flies

Mix 4 tablespoons (2 fluid oz.) of KILL-KO FLY SPRAY CONCENTRATE per 1 1/2 pints of water (1% concentration). Using a calibrated sprayer, apply 5 ml of spray to the face and a total of 5-10 ml to the back and sides of the animal 3 times per week. Improved control may be achieved by the addition of 1 level teaspoon of sugar per pint of spray solution.

For Control of Lice and Winter Ticks on Dairy Cattle, Beef Cattle, Sheep, Goats and Hogs.

Mix 1/2 pint of KILL-KO FLY SPRAY CONCENTRATE per 3 1/5 gallons of water (0.25% concentrate). Spray animals thoroughly using up to 1 gallon of spray. Apply a second application 14 days later. Repeat applications as necessary to maintain control, but not more often than once every 7 days.

Backrubber Applications

For use in backrubbers or face rubbers for the control of horn flies and face flies, mix 1 pint of KILL-KO FLY SPRAY CONCENTRATE per 2 gallons of backrubber base oil (1% concentrate). This may be diesel fuel or fuel oil in accordance with local recommendations.

1175-759

EPA Reg. No. 70-155

EPA EST. NO. KY 1

Manufactured by
RIGO COMPANY
Buckner, Kentucky 40010



FLY SPRAY CONCENTRATE

EMULSIFIABLE CONCENTRATE DAIRY AND LIVESTOCK SPRAY

Contains CIODRIN® and VAPONA® Insecticides

ACTIVE INGREDIENTS	BY WEIGHT
Dimethyl phosphorothioate of Alpha-methyl benzyl 3-hydroxy-2-naphthyl	10.0%
2,2-bis(4-chlorophenyl) ethyl phosphate	2.3%
Phenyl Glycidyl Ether	0.2%
Heavy Aromatic Hydrocarbons	77.5%
INERT INGREDIENTS	10.0%
TOTAL	100.0

*Equivalent to 2% Vapona® Insecticide
CIODRIN® and VAPONA® are
Registered SHELL® Trademarks

NET CONTENTS: ONE QUART

WARNING: Keep Out of Reach of Children

See side panel for additional cautions

DIRECTIONS FOR LIVESTOCK PEST CONTROL

For Control of House Flies, Face Flies, and Stable Flies In and Around Livestock Buildings.

Mix 1 pint of KILL-KO FLY SPRAY CONCENTRATE per 1 1/2 gallons of water (1% concentration). Spray premise such as inside and outside barn walls (including dairy barns), eaves, fences and other resting or breeding places at least twice monthly using approximately 1 gallon of finished spray per 1000 square feet (40' x 25'). Improved house fly and face fly control may be achieved by adding sugar to the spray solution at the rate of 1 pound per 6-10 gallons of spray.

PRECAUTIONS IN USING

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS
WARNING: HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. DO NOT GET IN EYES, ON SKIN OR UN CLOTHING. DO NOT BREATHE SPRAY MIST, KEEP AWAY FROM HEAT AND OPEN FLAME

Wash thoroughly with soap and water after handling and before eating and smoking. Wear clean clothing. In case of spilling on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

During commercial or prolonged exposure in spray-mixing and loading operations, wear clean rubber gloves and a mask or respirator of a type found suitable by the U.S. Bureau of Mines for protection against KILL-KO FLY SPRAY CONCENTRATE.

Do not apply externally in combination with Trichloroethene. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place. This container is not for use or storage in or around the home.

Do not contaminate feed or foodstuffs. Do not use in poultry houses. Do not treat animals within one day of slaughter.

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.
NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

BEST AVAILABLE COPY

in accordance with
for Phase 2
Change in Status.

A. ACTIVE INGREDIENTS:

Dimethyl phosphate of Alpha-methylbenzyl 3-hydroxy-
cis-crotonate10.0%
Dichlorvos [2, 2-dichlorovinyl dimethyl phosphate]
.....2.3%*
Related compounds.....0.2%*

INERT INGREDIENTS**:.....87.5%
TOTAL 100.0%

*Equivalent 2.5% Vapona^(R) Insecticide.

**Contains Heavy Aromatic Naphtha.

B. Rigo Company

A Division of Wilbur-Ellis Company
Buckner, KY 40010