

RIGO'S BEST

For use on fruit trees, and ornamental trees and shrubs. Controls borers and other listed insects.

LINDANE
INSECTICIDE CONCENTRATE

A

ACTIVE INGREDIENTS:

Lindane (Gamma Isomer of Benzene Hexachloride)....	20.0%
Naphtha.....	38.2%
INERT INGREDIENTS.....	41.8%
TOTAL.....	100.0%

EPA Reg. No. 70-163

EPA Est. No. 70-KY-1

CAUTION Keep Out Of Reach Of Children

See adjoining panel for additional precautionary statements

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Keep out of reach of children. Children should not be allowed to handle or apply this product. Harmful if swallowed. Avoid inhalation of spray mist. Avoid skin contact. Harmful vapors may be absorbed through the skin. Wash thoroughly with soap and water after using and before eating or smoking. Avoid contamination of food or feedstuffs. Food utensils such as teaspoon or measuring cup should not be used for food purposes after using with insecticide.

STATEMENT OF PRACTICAL TREATMENT

If swallowed - Call physician or Poison Control Center immediately. If possible vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled - Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin - Remove contaminated clothing and wash affected areas with soap and water.

If in eyes - Flush eyes with plenty of water. Call a physician immediately.

NOTE TO PHYSICIAN - Solvent presents aspiration hazard. Gastric lavage is indicated if material was taken internally.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, birds, and other wildlife. Do not apply directly to water or wetland. Drift and runoff from treated areas may be hazardous to fish in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

This pesticide is highly toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually between late evening

29 3 5

and early morning.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill, spray, or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Store in original container in a cool, dry location away from heat, flame, direct or indirect sunlight. Do not reuse empty container. Wrap container and put in trash collection.

NOTE: Read precautionary statements and directions thoroughly before using. Apply this product only as specified on the label. Avoid exposure to children. Do not allow children to handle or apply this product. Do not allow children or pets in treated areas until surfaces are dry.

B → NET CONTENTS 1 QUART
MANUFACTURED BY RIGO CO.,
BUCKNER, KY. 40010

For complete use directions see attached brochure

**RIGO'S BEST
LINDANE**

INSECTICIDE CONCENTRATE

For use on fruit trees, and ornamental trees and shrubs. Controls borers and other listed insects.

SEE CONTAINER LABEL FOR INGREDIENT STATEMENT, COMPLETE PRECAUTIONARY STATEMENTS INCLUDING STATEMENT OF PRACTICAL TREATMENT.

READ ENTIRE LABEL

TREES AND SHRUBS - For control of borers and other listed insects.

PROTECTIVE CLOTHING: Applicators must wear long sleeved shirt, long pants, waterproof gloves, full foot covering (such as shoes or boots), and head covering (such as a hat).

MIXING and APPLICATION: See table below.

Borer and Other Insects	Trees or Shrubs	Dosage	Time to Apply and Remarks
Leaf Miners	Elm, Locust, Birch	1 tsp/ 1 gal. water	Apply when small miners appear in early and mid-May. For Locust early July.
Twig Girdler	Junipers	1 tbl/ 1 gal. water	Make 2 applications at 30 day intervals. No. Calif. - June and

Lindane Ornamental (labels\lindanor)
PAGE 3

Mid-July. So. Calif.
April and May.

Pine Bark Beetles	Pines	1/2 pt./ 2 1/2 gals. 1 gal. water	Spray trunks thoroughly - wet trunks to run-off.
White Pine Weevil	Pine, Spruce	1 tbls/ gal. water	Apply late April when buds begin to swell. Spray upper parts of trees thoroughly.
Taxus Weevil	Taxus, Hemlock	1 tbls/ 1 gal. water	Apply in May or early June to the trunks and around base of trees soaking the barks and wetting the soil surface.
Bark Beetles, Wood Boring Beetles	Maple, Aspen, Poplar, Ash, Willows, Elm Locust	2 tbls/ 1 gal. water	Apply in spring when buds begin to break. Elms-During summer. Soak trunks to run- off.
Thrips, Plum Curculio, Flea Beetles	Apples, Pears, Plums, Peaches Cherries	1 teas/ 1 gal. water	Do not use on Apple, Pear, or Plum trees within 60 days of harvest. Do not apply to Peaches later than seven days after shuck split.
Fruit Tree Bark Beetles, Flat-Headed Borers, Round-Headed Borers	Apples Cherries Peaches	1 tbls/ 1 gal. water	Apply to bark of tree trunk and large limbs in spring. The appli- cation limitation in the label directions for the listed fruit crops should be observed. Apply to trunk, crotch to ground. Soak bark to run-off. Do not apply to fruit or foliage.
Peach Tree Borer	Peaches	1 tbls/ 1 gal. water	Spray lower trunk and around base of tree, soaking bark and

4 of 5

wetting trunk from
crotch to ground. Do
not apply to fruit or
foliage.

Dogwood Borer, Rhododendron Borer, Lilac Borer	Dogwood, Rhododendron, Lilac	1 tbs/ 1 gal. water	Make 2 to 3 appli- cations 7 to 10 days apart in May and early June.
--	------------------------------------	---------------------------	---

Iris Borer	Iris	1 tsp/ 1 gal. water	In early spring, des- troy old leaves. Apply during April and May.
------------	------	---------------------------	---

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

B →

Manufactured by RIGO COMPANY, BUCKNER, KENTUCKY 40010

Attachment (70-163)

A. ACTIVE INGREDIENTS:	
Lindane (Gamma Isomer of Benzene Hexachloride).	20.0%
INERT INGREDIENTS*:	80.0%
TOTAL	100.0%

*Contains Heavy Aromatic Naphtha.

B. Rigo Company
 A Division of Wilbur-Ellis Company
 Buckner, KY 40010