

**DIRECTIONS
CONTINUED**

moth. Use 5 lbs. per 5,000 square feet. To control — Plum curculio and codling moths apply at petal fall and every 10 to 14 days thereafter as long as local conditions warrant.

GRAPES: To control Grape leaf folder and Grape Berry moth and Japanese beetle, apply the dust to thoroughly cover upper and lower leaf surfaces. Apply just before first brood larvae emerge from rolls. Apply as needed for leafhoppers.

STRAWBERRIES: To control—Meadow spittlebug and Strawberry leaf roller apply the dust to thoroughly cover upper and lower leaf surfaces.

FLOWERS, SHRUBS, ORNAMENTAL TREES: To control — Apple aphid, Bag-worms, Birch leaf miner, Blister beetles, Boxelder bugs, Boxwood leaf miner, Elm leaf aphid, Elm leaf beetle, Flea beetles, Japanese beetles, Lace bugs, Leafhoppers, leaf rollers, Oak leaf miner, Orange tortrix, Periodical cicada, Plant bugs, Psyllids, Rose aphids, Rose-slug, Scale insects, Tent caterpillars, Willow leaf beetle and exposed Thrips, apply the dust to thoroughly cover upper and lower leaf surfaces. Apply when insects or damage first appears and repeat at weekly intervals or as needed. For scale insect control, apply in spring and early summer when scale crawlers are present.

TOBACCO: To control—Hornworms, Budworms and Flea beetles, apply the dust to thoroughly cover upper and lower leaf surfaces. Where budworm damage is occurring, apply to entire plant, directing the dust downward into the bud; repeat as necessary. Treated fields may be entered immediately after dust has settled. If late applications are necessary, treat immediately after priming rather than before priming. Avoid heavy applications as plant injury may occur.

POULTRY: Poultry Houses: For use in and around poultry houses. To control Chicken mites, Fleas, Bed bugs, Fowl ticks, use one half pound per 40 sq. ft. of wall, bedding, litter or roosts surfaces. Force dust into cracks and crevices. Ventilate while dusting. Repeat as needed. Do not treat nest litter. Avoid direct contamination of eggs. Do not use on poultry or game birds. Do not use in premise treatment within 7 days of slaughter.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label in-

KILL-KO KILL-KO KILL-KO KILL-KO KILL-KO KILL-KO KILL-KO KILL-KO KILL-KO KILL-KO

KILL-KO.

10% SEVIN*
DUST

ACTIVE INGREDIENTS:
CARBARYL (1-Naphthyl N-methylcarbamate)10.00%
INERT INGREDIENTS:90.00%
*Trademark of Union Carbide Chemical Company

CAUTION
KEEP OUT OF REACH OF CHILDREN
See back panel for additional caution.

MANUFACTURED BY
RIGO CHEMICAL COMPANY
NASHVILLE, TENN. 37202

