

70-165 7-30-2003 1-4

Read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States Environmental Protection Agency Washington, DC 20460

Registration Amendment Other (checked)

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number: 70-165
2. EPA Product Manager: Insecticide-Rodenticide Branch
3. Proposed Classification: None (checked), Restricted
4. Company/Product (Name): Kill-Ko 10% Sevin brand carbarvl insecticide Dust
5. Name and Address of Applicant: Value Garden Supply, LLC, P. O. Box 585, St. Joseph, MO 64502
6. Expedited Review: In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. Product Name

Section - II

Amendment - Explain below. Resubmission in response to Agency letter dated. Notification - Explain below. Final printed labels in response to Agency letter dated. "Me Too" Application. Other - Explain below. NOTIFICATION JUL 30 2003

Explanation: Use additional page(s) if necessary. (For section I and Section II.) Notification of label revision updating disposal instructions in response to PR Notice 2001-6; This notification is consistent with the guidance in PR Notice 2001-6 and the requirement of EPA's regulations at 40 CFR 156.10 and 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product.

Section - III

1. Material This Product Will Be Packaged In: Child-Resistant Packaging, Unit Packaging, Water Soluble Packaging, 2. Type of Container: Metal, Plastic, Glass, Paper, Other (Specify)
3. Location of Net Contents Information: Label, Container
4. Size(s) Retail Container
5. Location of Label Directions
6. Manner in Which Label is Affixed to Product: Lithograph, Paper glued, Stenciled, Other

Section - IV

1. Contact Point: Name: Carrie Crawford, Title: Regulatory Affairs Manager, Telephone No.: 540-864-8455
Certification: I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete.
2. Signature: Carrie Crawford
3. Title: Regulatory Affairs Manager
4. Typed Name: Carrie Crawford
5. Date: July 14, 2003
6. Date Application Received (Stamped):

FOR HOMEOWNER USE ONLY

2-4

## KILL-KO 10% SEVIN® brand carbaryl insecticide DUST

### ACTIVE INGREDIENTS

Carbaryl: (1-Naphthyl N-methylcarbamate).....	10%
Other Ingredients:.....	90%
TOTAL	100%

Sevin® is a registered trademark of Rhone Poulenc for carbaryl insecticide.

EPA Reg. No. 70-165  
EPA Est. No. 44515-MO-1

Distributed by: Value Garden Supply, LLC.  
P. O. Box 585  
St. Joseph, MO 64502  
(952) 884-6477  
www.valuegardens.com

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See Side/Back Panel for Additional Precautionary Statements

NET WEIGHT:

### FIRST AID

- If swallowed:**
- Call poison control center or doctor immediately for treatment advice.
  - Have person sip a glass of water if able to swallow.
  - Do not induce vomiting unless told to do so by the poison control center or doctor.
  - Do not give anything by mouth to an unconscious person.
- If inhaled:**
- Move person to fresh air.
  - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
  - Call a poison control center or doctor for further treatment advice.
- If on skin or clothing:**
- Take off contaminated clothing.
  - Rinse skin immediately with plenty of water for 15-20 minutes.
  - Call a poison control center or doctor for treatment advice.
- If in eyes:**
- Hold eye open and rinse slowly and gently with water for 15 – 20 minutes.
  - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
  - Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Note to Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

For information of this product (including health concerns, medical emergencies or pesticide incidents) call the National Pesticide Information Center at 1-800-858-7378.

### PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

**CAUTION:** Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling.

### ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**USER SAFETY REQUIREMENTS**

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until dust has settled.

Wear household latex or rubber gloves; use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust.

Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using a liquid carbaryl product. Do not repeat application more than once a week.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**VEGETABLES:** Use only the amount of dust necessary to provide adequate coverage. Begin applications when insects or damage appears and continue at 7 to 10 day intervals, as needed, for control. The pre-harvest interval indicates the number of days between the last application and harvest.

**BEANS:** To control Mexican Bean Beetle, apply the dust to thoroughly cover upper and lower leaf surfaces. Also to control Bean Leaf Beetle, Cucumber Beetle, Western Bean Cutworm, Flea Beetle and Japanese Beetle, Corn Earworm, Armyworms and Velvetbean Caterpillar, Lima Bean Pod Borer, Lygus Bug and Stink Bug, apply the dust to thoroughly cover upper and lower leaf surfaces. For Beans, Black-Eyed Peas, Cowpeas, Crowder, Southern Peas, Dry Beans, Green Beans, Lima Beans, Navy Beans and Snap Beans a 3-day pre-harvest interval is required for succulents and 21 days for dried.

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI:** To control Flea Beetle, Harlequin Bugs, Corn Earworm, Armyworm, Cabbage Caterpillars, and small Cabbage Loopers, apply the dust thoroughly to cover upper and lower leaf surfaces. A 3-day pre-harvest interval is required.

**CORN:** To control Corn Earworms, European Corn Borer, Fall Armyworms, Sap Beetles and Japanese Beetles, apply the dust to thoroughly cover upper and lower leaf surfaces. For Corn Earworm begin when first silks appear and direct to the zone of ear production. Apply until the silks begin to dry but not more often than once a week.

**CAUTION:** Application of this product to the tassel region of the corn during pollen shed period will seriously reduce bee populations. Pre-harvest Interval - 2 days for sweet corn ears; 14 days for forage and 48 days for fodder. Field and popcorn is 14 days for forage and silage and 48 days for grain and fodder.

**TOMATO, EGGPLANT, PEPPER:** To control Tomato Fruitworm, Fall Armyworm, Flea Beetle and Lygus Bugs, apply dust to thoroughly cover upper and lower leaf surfaces. A 3-day pre-harvest interval is required.

**CUCUMBER, SQUASH, CANTALOUPE, PUMPKINS, MELONS:** To control Pickleworm, Melonworm, Cucumber Beetle, Squash Bug, Flea Beetles, apply dust to thoroughly cover upper and lower leaf surfaces. A 3-day pre-harvest interval is required.

**LETTUCE:** To control Flea Beetles, Harlequin Bug, Leafhopper, Corn Earworm, Armyworm, Cabbage Caterpillars, Spittlebugs, Lygus Bugs and Stink Bug, apply dust to thoroughly cover upper and lower leaf surfaces. On head lettuce, do not apply within 14 days of harvest. On leaf lettuce, a 14-day pre-harvest interval is required.

**CARROTS:** To control Flea Beetles, Leafhoppers, Spittlebugs and Stink Bug, apply dust to thoroughly cover upper and lower leaf surfaces. A 7-day pre-harvest interval is required.

**OKRA:** To control Corn Earworm and Stink Bug, apply dust to thoroughly cover upper and lower leaf surfaces. A 3-day pre-harvest interval is required.

**POTATO:** To control Colorado Potato Beetle, Leafhopper, Flea Beetles, and Lygus Bug, apply the dust to thoroughly cover upper and lower leaf surfaces. A 7-day pre-harvest interval is required.

**FRUIT:** Use only the amount of dust necessary to provide adequate coverage. Begin applications when insects or damage appears and continue at 7 to 10 day intervals, as needed, for control. The pre-harvest interval indicates the number of days between the last application and harvest.

**STRAWBERRIES:** To control Meadow Spittle Bug and Strawberry Leaf Roller, apply dust to thoroughly cover upper and lower leaf surfaces. A 7-day pre-harvest interval is required.

4-4

**FLOWERS AND SHRUBS:** To control Flea Beetles, Japanese Beetles, Lace Bugs, Leafhoppers, Leaf Rollers, Plant Bugs, Psyllids, Rose Aphids, Rose Slug, Scale Insects, Tent Caterpillars, and Exposed Thrips, apply dust to thoroughly cover upper and lower leaf surfaces. Apply when insects or their damage first appears and repeat at weekly intervals if needed. For scale insect control, apply in spring and early summer when scale crawlers are present.

#### POULTRY

**DIRECT DUST APPLICATION TO BIRDS:** Use ½ pound per 100 birds. Apply by shaker can or hand duster for control of northern fowl mite and lice and as a supplement to premise treatment for chicken mites and fleas. Directly dust each bird and fluff feathers. **DO NOT APPLY WITHIN 7 DAYS OF SLAUGHTER.** Repeat treatment in 4 weeks if necessary. Direct dusting is a supplement to litter treatment for control of the above insects.

**LITTER TREATMENT:** Apply thoroughly to floors, roosts and interior surfaces, using ½ pounds per 40 square feet to control chicken mites, fleas and bed bugs. Apply by shaker can or hand duster. Do not apply directly to eggs or nest litter. Do not house birds in treated houses within 7 days of slaughter. Do not contaminate feed or drinking water.

**POULTRY DUSTING BOX:** For control of northern fowl mite, chicken mite, lice and fleas, place one self-treatment dust box (18x12x3) containing 1 ¼ pound 10% Sevin Dust per each 50 birds. Allow birds to dust themselves. This is a supplement to litter treatment for chicken red mite and flea control. Remove birds from dust boxes 7 days before slaughter.

#### DOGS AND CATS

Use this product only outside in a well ventilated area. Wearing household latex or rubber gloves, dust entire animal, avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. This product should not be applied by children. Do not use on pregnant dogs. Do not use on kittens or puppies under 12 weeks of age. Dust bedding and doghouse using a shaker can or other dust applicator.

**FOR FLEAS, LICE AND BROWN DOG TICKS:** Brush hair back and apply powder down to the skin and directly to ticks. Sprinkle powder on floor of kennel in cracks, crevices and other hiding places. If the animal has free run, treatments may have to be made every second or third week, but no more frequently than once a week. To avoid unnecessary applications, animal should be closely examined for fleas before retreating. If the animal is confined to an area free of fleas, one treatment may be sufficient. Repeat if necessary but no more frequently than once a week.

For control of fleas in kennels, pens, yards, lawns and under houses, remove manure or debris and dust approximately 4 oz. on an area up to 180 square feet.

#### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in a cool, dry area away from heat or open flame. Keep in original container and preferably in a locked storage area inaccessible to children and pets.

**Disposal:** If Empty: Do not reuse this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain

Buyer assumes all responsibility for safety and use not in accordance with label directions.