DIRECTIONS

BEFORE USE READ CAUTION AND NOTICE STATEMENTS CAREFULLY

SPRAY APPLICATION: [illegible] material will make 6¼ gallons of fruit tree spray. For [illegible] use 5 level tablespoonfuls to each gallon of water. Mix well. Agitate solution while spraying. Apply spray to trees thoroughly.

APPLES, PEARS: To control Bitter Rot, Scab, Brooks Fruit Spot, Woolly and Green Aphids, Pacific, Clover and Red Spider Mites, Codling Moth, Forbes Scale, Plum Curculio, Apple Maggot, Japanese Beetle, Red Banded and Leaf Roller, apply as follows:

1. When blossom stems separate and before blossoms open.
2. When 75% of the petals have fallen.
3. 10 days later.
4. 10 days to 2 weeks later.
5. Continue at 2-week intervals to 21 days before harvest.

Do not use on D'Anjou pears. Russetting may be produced on Bosc pears. Red delicious and other sensitive varieties may be injured by early season application.

PEACHES: To control Brown Rot, Peach Scab, European Red Mite, Red Spider, Plum Curculio, Oriental Fruit Moth, Japanese Beetle, [illegible] Fruit Scale, Pacific and Clover Mites, apply as follows:

1. Full bloom.
2. When shucks begin to split.
3. 10 days later.
4. 10 days later.
5. 2 weeks later.
6. Do not apply within 21 days of harvest.

PLUMS (PRUNES): To control Pacific, European and Red Spider Mites, Plum Curculio, Mealy Plum and Green Peach Aphids and Japanese Beetle, apply as follows:

1. When trees are in full bloom.
2. When shucks are falling.
3. 10 days later.
4. 2 weeks later.
5. Continue at 2 week intervals if necessary. Do not apply later than 21 days before harvest.

If [illegible] of diseases on above fruit crops is necessary within 21 days of harvest, use [illegible] spray. Remove residues by washing, [illegible].

CAUTION

Harmful if swallowed. Avoid inhalation of dust or spray mist. Avoid [illegible] and wash thoroughly after using. Do not [illegible] feed or feedstuffs. Keep out of [illegible] quantity, limit of method of application [illegible] property or crops or [illegible] treatment or [illegible] from areas treated [illegible].

Use of this product other than indicated on the label Buyer assumes all

Manufactured by
RIGO COMPANY
BUCKNER, KENTUCKY 40010

KILL-KO®

FRUIT TREE SPRAY

ACCEPTED
OCT 17 1974
UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. [illegible]

protects your fruit trees

RECEIVED
8 OCT 1974
REGISTRATION DIVISION
PESTICIDES, EPA

CONTAINS
Methoxychlor*
Malathion
Captan

An insecticide and fungicide for insect and disease control on the fruit trees

ACTIVE INGREDIENTS:

Technical Methoxychlor*	[illegible]
Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide)	[illegible]
Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate)	[illegible]
INERT INGREDIENTS	[illegible]
	[illegible]

CAUTION: Keep Out of the Reach of Children

See back panel for other cautions

[illegible] POUNDS NET WEIGHT

Manufactured by
RIGO COMPANY
BUCKNER, KENTUCKY 40010