

## DIRECTIONS FOR USE

For the control of the following tomato insects: Blister Beetles, Colorado Potato Beetle, Flea Beetles, Green Peach Aphids, Tomato Fruitworm, Yellow-striped Armyworm, Tomato Hornworm, Southern Green Stink Bug, White Fly. Applications should be started as soon as insect pests are first observed and unless otherwise directed. Should be repeated at 7 to 10 day intervals throughout the season.

For the control of the following tomato diseases: Early and late blight, gray leaf spot (stemphyllium), anthracnose, septoria leaf spots and cladosporium gray mold. For disease control, begin applications before diseases appear and repeat at 7 to 10 day intervals.

Apply this material in a manner that will give a uniform coating on all above ground surfaces of the plant. When dusting, be certain to cover both the lower and upper surface of foliage. During periods of extremely high temperatures, it is preferable to dust in the late evening or early morning.

Application may be made up to five (5) days before harvest.

**CAUTION:** Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed or foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.


This product is toxic to bees and should not be applied when bees are actively visiting the area. Highly toxic to fish and wildlife. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

Dispose of container when empty by burying.

2141-792


EPA Reg. No. 70-173

Manufactured by  
**RIGO Chemical Company**  
Nashville, Tenn. 37202



# TOMATO DUST

ACCEPTED  
OCT 11, 1972  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND ROENTHOGEN ACT  
FOR ECONOMIC POISON RESIST  
ED UNDER NO. 2-173  
SOPHOMORE  
COMMITTEES



Insecticide — Fungicide

**ACTIVE INGREDIENTS:**

\*Endosulfan (Hexachlorohexahydromethano-  
2,4,3-benzodioxathiepin oxide) 4.0%

\*\*Zineb 6.5%

**INERT INGREDIENTS**

\*Thiodan U. S. Pat. No. 2,799,685 89.5%  
\*\*Zinc ethylene bisdithiocarbamate  
Zinc expressed as metallic 1.54%  
Thiodan is a registered trademark of  
Farbwerke Hoechst A.G.

NET WEIGHT 1 POUND

**CAUTION**  
KEEP OUT OF THE REACH OF CHILDREN  
See back panel for additional caution.

Manufactured by  
**RIGO Chemical Company**  
Nashville, Tenn. 37202