


DIRECTIONS



Apply Kill-Ko SUGAR FLY BAIT daily while flies are abundant. Sprinkle thinly in strips on floor and other places where flies gather, but not where it will contaminate feed or food. Usually, about 1 oz. (2 tbsp.) is needed for each 500 sq. ft., depending on the number of flies. If they are extremely numerous, several times this quantity may be necessary.

Kill-Ko SUGAR FLY BAIT is especially recommended for use in barns, stables, pens, feed rooms, poultry houses, kennels, manure piles, garbage cans, and other outside areas. In dairy barns, treat the feed room floor and the barn floor, but do not feed the floor sweepings to livestock. Repeat as needed.

In poultry houses where poultry is in cages, use SUGAR FLY BAIT on walkways in and around the building, including feed rooms and egg houses. Where poultry is on ground or litter, apply only outside the building or in yards where poultry cannot feed on Fly Bait.


To kill fly maggots, apply Kill-Ko SUGAR FLY BAIT evenly and thoroughly on manure at the rate of 1 to 1½ lbs. per 100 sq. ft. (2 to 2½ oz. per sq. yd.) If in small mounds, apply on sides as well as top. Repeat in 5 to 7 days, and afterwards as needed.

KEEP TIGHTLY CLOSED WHEN NOT IN USE

Manufactured by

RIGO CHEMICAL COMPANY • NASHVILLE, TENN. 37203

SUGAR FLY BAIT



VAPONA®

**Attracts
and kills
RESISTANT
House Flies**

VAPONA is a registered Shell trademark

ACTIVE INGREDIENTS	2, 2-Dichlorovinyl dimethyl phosphate	0.44%
	Rotenone Compounds	0.04%
INERT INGREDIENTS	99.52%

NET WEIGHT: 1 LB.

EPA Reg. No. 70-1195

CAUTION: KEEP OUT OF REACH OF CHILDREN. See side panel for additional caution.

Harmful if swallowed or absorbed through skin. Avoid skin contact. Wash thoroughly with soap and water after using. Keep container out of reach of children and animals. Avoid contamination of food, feed, and utensils used for food or feed. Do not use inside homes or where milk and other foods are stored or processed. Repeated use may cause Cholinesterase inhibition. Atropine is antidotal.

Food utensils such as teaspoons and table-spoons should not be used for food purposes after use with pesticides. Bait should be placed in areas inaccessible to children and pets.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and/or handling is contrary to label instructions.

BAIT

CTIONS

...R FLY BAIT dairy while flies
...thly in strips on floor and
...gather, but not where it will
...Usually about 1 oz (2 tbs) p
...ft, depending on the num-
...extremely numerous, several
...be necessary.

...LY BAIT is especially recom-
...stables, pens, feed rooms,
...manure piles, garbage cans.
...In dairy barns treat the feed
...floor, but do not feed the floor.
...Repeat as needed.

...re poultry is in cages, use
...BAIT on walkways in and
...cluding feed rooms and egg
...s on ground or litter, apply
...p or in yards where poultry.

...ply ... SUGAR FLY
...hly on manure at the rate of
...ft (2 to 2 1/2 oz per sq yd)
...ly on sides as well as top
...d afterwards as needed.

NOT IN USE

KILL-KO®

SUGAR FLY BAIT



VAPONA®

**Attracts
and kills
RESISTANT
House Flies**

VAPONA® is a registered Shell trademark.

ACTIVE INGREDIENTS	2,2-Dichlorovinyl dimethyl phosphate	0.46%
	Related Compounds	0.04%
INERT INGREDIENTS		99.50%

NET WEIGHT: 1 LB.

EPA Reg. NO. 70-176

CAUTION: KEEP OUT OF REACH OF CHILDREN. See side panel for additional caution.

READY TO USE

CAUTION

Harmful if swallowed or absorbed through skin. Avoid skin contact. Wash thoroughly with soap and water after using. Keep container out of reach of children and animals. Avoid contamination of food, feed, and utensils used for food or feed. Do not use inside homes or where other foods are stored or processed. Repeated use may cause Cholinesterase inhibition. Atropine is antidotal.

Food utensils such as teaspoons and spoons should not be used for food preparation after use with pesticides. Bait should be used in areas inaccessible to children and pets.

This product is toxic to fish and wildlife. Feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy v

NOTICE: Seller makes no warranty, expressed or implied, concerning the quality of this product or its use. Buyer assumes all risk of use and or handling of this material when use and or handling is contrary to label directions.