

RIGO'S BEST DIAZINON 2E

ACCEPTED
NOV 15 1993
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 70-177

**KILLS OVER 100 DESTRUCTIVE:
LAWN & GARDEN PESTS**

Lawn & Garden Insect Control

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

Active Ingredient:
 Diazinon: O, O-Diethyl
 O-(2-Isopropyl-6-
 methyl-4-pyrimidinyl)
 phosphoro-
 thioate 25.0%

Inert
 Ingredients*: 75.0%

Total: 100.0%

* Contains aromatic petroleum distillates.

**KEEP OUT OF REACH
OF CHILDREN.**

CAUTION

See side panel for additional precautionary statements.

EPA Reg. No. 70-177

Protects
 Lawns
 Vegetables
 Fruits and Nuts
 Roses, Flowers
 Trees, Shrubs

Controls
 Ants, Aphids
 Chinch Bugs
 Cutworms
 Fire Ants¹
 Flea Beetles
 Grubs, Mites
 Sod Webworms.

...and many others

¹Aids in control

See full label in booklet under cap for complete directions for mixing and applying this product and a complete list of pests controlled.

One Pint
Net Contents

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

General Information

Diazinon 2E Insect Spray is an emulsifiable concentrate product which must be diluted with water before application. This product kills many outdoor pests which damage lawns, roses, flowers, trees, shrubs, vegetables and fruits and insects which annoy pets and people.

SEE THE BOOKLET UNDER CAP FOR COMPLETE DIRECTIONS FOR USE AND PESTS CONTROLLED.

Notes: (1) USE CONTENTS WITHIN 6 MONTHS AFTER OPENING. SEE STORAGE AND DISPOSAL DIRECTIONS. (2) Do not use in the home. (3) Do not use on ferns, pansy, hibiscus, papaya, olea, geranium or orchids since plant injury may occur. (4) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (5) Do not apply to food crops grown in greenhouses.

After spraying, wait the following number of days before harvesting fruits, nuts, or vegetables:

1 day	Tomatoes
3 days	Melons, Winter Squash
5 days	Peppers, Strawberries
7 days	Beans, Broccoli, Blackberries, Boysenberries, Cauliflower, Cucumbers, Dewberries, Loganberries, Peas, Raspberries, Summer Squash, Sweet Corn
10 days	Chinese Broccoli, Chinese Cabbage, Collards, Kale
14 days	Carrots, Endive (Escarole), Lettuce, Onions (Bulb and Green), Parsnips, Radishes, Spinach, Turnips
21 days	Apples, Cabbage, Cherries, Nectarines, Peaches, Plums, Prunes
28 days	Graes
35 days	Potatoes (Irish)
45 days	Walnuts
*	Almonds (one dormant spray only)

Precautions: (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water and launder clothing before wearing again.

* This table is only a partial listing of the pests controlled by this product. See booklet under cap for complete directions for use and pests controlled.

Problem Area	Pests*
Nuisance Pests in Outside Areas	Ants, Brown Dog Ticks, Chiggers, Cockroaches, Crickets, Fleas, Fire Ants, Soiders and others*
Lawns	Chinch Bugs, Ants, Sod Webworms, Brown Dog Ticks, Fleas, White Grubs, Lawn Billbugs and others*
Vegetables: Foliar Application	
Beans, Peas (except dried varieties)	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites
Broccoli, Chinese Broccoli, Cabbage, Chinese Cabbage, Cauliflower, Collards, Kale	Apples, Flea Beetles, Dipterous Leafminers, Diamondback Moths, Imported Cabbageworms
Endive (Escarole), Lettuce, Spinach	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites

BEST AVAILABLE COPY

Peppers. Tomatoes	Aonids. Colorado Potato Beetles. Fall Armyworms (Tomatoes). Flea Beetles. Leathoppers. Southern Armyworms (Tomatoes). Vinegar Flies - <i>Drosophila</i> sp. (Tomatoes)
Onions (Bulb and Green)	Onion Thrips
Radishes. Turnips. Carrots. Parsnips. Potatoes	Aonids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Colorado Potato Beetles (Potatoes). Southern Armyworms (Potatoes)
Sweet Corn	Aonids. Flea Beetles. Grasshoppers. Spider Mites
Melons. Squash (Summer, Winter)	Aonids. Thrips. Spider Mites
Vegetables: Preplant Soil Application	
Beans. Peas (except dried varieties)	Cutworms. Mole Crickets. Wireworms (Snap Beans)
Beets. Red (Table)	Cutworms. Mole Crickets
Cabbage	Cutworms. Root Maggots. Wireworms
Carrots. Sweet Corn	Cutworms. Mole Crickets. Wireworms
Lettuce. Tomatoes	Cutworms. Wireworms
Radishes. Turnips	Cutworms
Fruits & Nuts	
Almonds (California only)	Scale Crawlers
Walnuts (California only)	Aonids. Scale Crawlers. Spider Mites and others *
Apricots. Peaches. Nectarines	Aonids. Apricot Mealybugs (Apricots). Spider Mites. Scale Crawlers
Cherries	Black Cherry Aonids. Cherry Fruitflies. Cherry Rust Mites. Scale Crawlers and others *
Plums. Prunes	Aonids. Spider Mites
Caneberries (CA, OR, WA only): Blackberries. Boysenberries. Dewberries. Loganberries. Raspberries	Aonids. Leafhoppers. Two-Spotted Spider Mites. Thrips. Raspberry Fruitworms
Strawberries	Aonids. Two-Spotted Spider Mites. Strawberry Leafrollers
Grapes	Aonids. Grape Berry Moths. Grape Leafrollers. Leafhoppers
Roses & Flowers	
Azaleas. Camellias. Chrysanthemums. Carnations. Gladiol. Lilac. Rhododendron. Roses	Aonids. Flea Beetles. Dipterous Leafminers. Leafhoppers. Mealybugs. Two-spotted Spider Mites. Thrips. Whiteflies and others *

BEST AVAILABLE COPY

Ornamental Trees & Shrubs

Arborvitae.

Aphids, Mites, Scale Crawlers, Whiteflies, Tent Caterpillars and others*

Azaleas, Birch.

Burwood, Camellias.

Douglas Fir, Elm.

Hawthorn Holly.

Juniper, Lilac.

Locust, Magnolia.

Mimosa, Oak, Pine.

Ornamental Plum.

Podar.

Rhododendron.

Spruce, Willow

Storage and Disposal

Storage

Do not store near heat or open flame. Store in original container in an area inaccessible to children or pets. **USE CONTENTS WITHIN 6 MONTHS AFTER OPENING.**

Disposal

Do not reuse empty container. Thoroughly rinse container, securely wrap in several layers of newspaper, and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on nontoxic pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

Statement of Practical Treatment

If swallowed: Call a physician immediately. **DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE.** Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention.

If on skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If in eyes: Rinse eyes with plenty of water and call a physician immediately.

Note to Physicians: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains aromatic petroleum

distillates which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply at least 1/4 inch of water immediately after spraying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water.

Spill may be hazardous to aquatic organisms in neighboring areas. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

59 13

Manufactured by
RIGO COMPANY,
Buckner, KY 40010

6.

6.

RIGO'S BEST DIAZINON 2E

KILLS OVER 100 DESTRUCTIVE
LAWN & GARDEN PESTS

ACCEPTED
NOV 15 1993
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 70-177

Lawn & Garden Insect Control

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

Active Ingredient:
Diazinon: O, O-Diethyl
O-(2-isopropyl-6-methyl-4-pyrimidinyl)
phosphorothioate 25.0%
Inert
Ingredients*: 75.0%
Total: 100.0%

* Contains aromatic petroleum distillates.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See end of booklet for additional precautionary statements

EPA Reg. No. 70-177

Protects

- Lawns
- Vegetables
- Fruits and Nuts
- Roses, Flowers
- Trees, Shrubs

Controls

- Ants, Aphids
- Chinch Bugs
- Cutworms
- Fire Ants¹
- Flea Beetles
- Grubs, Mites
- Sod Webworms
- ...and many others

¹Aids in control

One Pint
Net Contents

BEST AVAILABLE COPY

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

loading of spray. Where emulsion (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment, respect local treatment, water down with at least 1/4 inch of water immediately after spraying and product; however, avoid watering before puddling occurs.

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Menon and St. Augustinegrass. To control pests listed below, apply recommended rates.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

General Information

Diazinon DE Insect Spray is an emulsifiable concentrate product which must be diluted with water before application. Refer to the section Here to Mix and Apply. This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables, fruit and nuts, and insects which annoy pets and people. Contains diazinon insecticide which provides positive control of many "here-to-kill" sucking and chewing insects. Green worms that are applications should be repeated only as directed to maintain control. *Diazinon*

Notes: (1) USE CONTENTS WITHIN 6 MONTHS AFTER OPENING. SEE STORAGE AND DISPOSAL DIRECTIONS. (2) Do not use in the home. (3) Do not use on ferns, perennials, nodules, peonies, peas, gardens or orchids since injury may occur. (4) Use this product on succulent peas and beans only. Do not use on dried peas or bean varieties. (5) Do not apply to food crops grown in green houses.

Precautions: (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Soak in the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

How to Mix and Apply

Step #1: Determine ease of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Applications should be repeated only as directed to maintain control.

Step #2: Use clean sprayer. Carefully measure and mix the amount of D-2-n and water as suggested in the rate chart.

Step #3: Spray as directed in rate chart. Thorough coverage is important. Flush sprayer with clean water after each use.

See chart for complete list of use sites, crops, and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.

Measurement Conversions

- 1 tablespoon = 3 teaspoons
- 1 fl. oz. = 6 teaspoons or 2 tablespoons
- 6 fl. oz. = 3/4 cup
- 1 1/2 tablespoons in 3 gals. of water = 1 1/2 teaspoons in 1 gal. of water
- 2 tablespoons in 3 gals. of water = 2 teaspoons in 1 gal. of water

Lawn Pest Control

Precautions: Birds, especially waterfowl, feeding or resting on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and 1/2 over 30. Do not exceed maximum permitted label rates. Birds that have been recommended significantly increase potential hazards to birds and waterfowl. Avoid over-

Pests	Rate		Remarks
	<i>Diazinon DE Insect Spray</i>	Water	
Lawn Chinch Bugs	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	Thoroughly wet down grass a few hours before applying. In the North, if Chinch Bugs are the only problem, a lower rate of application can be used: 3 teaspoons in 3 gals. of water covers 125 sq. ft. or 4 fl. oz. in 24 gals. of water per 1,000 sq. ft.
Ants, Armyworms, Bermudagrass, Mite, Clover, Mite, Songspas (Collared), Crickets, Curworn, Digger Weeds, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sawbugs (Pillbugs)	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	Thoroughly wet down grass a few hours before applying. For ants and digger weeds, spot spray ant hills and weed ground near openings. For digger weeds, apply at dusk, when weeds are not active. Billbug control: Apply when activity is first observed or when chews or brown grass is present indicating damage from the insect. For frit flies, endow grass and water thoroughly before treatment, but delay application until grass is dry.
Brown Dog Ticks, Chiggers, Fleas	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	NOTE: Do not apply this product to animals. For brown dog ticks, spray grass and in shelter; particularly near house.
Hyperodes (Weevil)	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	Treat problem areas in mid-July and again in mid-Aug. Water grass thoroughly after application.
Milipedes	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	Thoroughly wet grass a few hours before applying.
Rhodesgrass, Scales	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	Apply when crawlers first emerge.
White Grubs of Japanese Beetle, European Chafer, Southern Chafer	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	Apply any time between late July and early October. Water grass thoroughly after application.

BEST AVAILABLE COPY

Mulch and Pests Outside the Home

<p>Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs, Pillbugs, Squirrels, Springtails, Collembola</p>	<p>Mix 6 fl. oz. in 3 gals. of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entrance ways, walls, garbage cans, tree trunks, and tree cracks and other places where insects can hide.</p> <p>To prevent entry into houses, spray a five-foot band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet. Repeat application as necessary.</p> <p>Do not treat animals with this product. Do not treat plants (either shrub tree trunks) at this mulch and pest rate.</p>
--	--

<p>Fire Ants</p>	<p>To aid in the control of fire ants in home lawns and other outdoor home recreation areas, apply 2 tablespoons in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness.</p> <p>NOTE: For best results, apply in cool weather 65-80°F. or in early morning or late evening hours. Treat new mounds as they appear.</p>
------------------	---

Insect Control on Ornamental Trees, Shrubs and Flowers

Trees and Shrubs	Insects	Rate per 3 Gals. of Water	Timing/Special Directions
<p>Arbutus, Azaleas, Birch, Jasmine, Camellias, Douglas Fir, Elm, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poinsettia, Rhododendron, Sorbus, Willow</p>	<p>Aphids, Bagworms, European Pine Shoot Moths, Holly Bud Moths, Dicotyledonous Leafminers (Arbutus), Azalea, Birch, Boxwood, Holly, Oak, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider, Opacus Root Weevil, Omnivorous Leaf-miners, Scale Crawlers (Cottony-cushion), Eucynimus, Lecanium, Pine Needle, Pitt, San Jose, Soot)</p>	<p>2 table- spoons</p>	<p>Apply as a thorough foliar spray. Wet leaves and branches to the drooping point. Try to penetrate dense foliage with spray.</p>
	<p>Webworms (Juniper, Mimosa), Whiteflies</p>	<p>2 table- spoons</p>	<p>Apply as a thorough foliar spray. Mimosa webworms: apply as a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals.</p>
	<p>Apple-and-Thorn Skeletonizers, Hemlock Chermes, Oak Looper, Obliquebanded Leafroller, Pear Slug, Tent Caterpillar, Webworms (Cottony-cushion, Fall)</p>	<p>2 table- spoons</p>	<p>Apply as a thorough foliar spray.</p>

Roses and Flowers	Insects	Rate per 3 Gals. of Water	Timing/Special Directions
<p>Azaleas, Camellias, Chrysanthemums, Carnations, Gladiolus, Lilac, Rhododendron, Roses</p>	<p>Aphids, Eucynimus, Scale Crawlers, Flea Beetles, Dicotyledonous Leafminers, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Two-spotted Spider, Thrips, Whiteflies)</p>	<p>2 table- spoons</p>	<p>Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage.</p>

Fruits and Nuts

Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted. Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.

Crop	PHI*
Almonds	**
Apricots	21
Blackberries	7
Boysenberries	7
Cherries	21
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Plums	21
Prunes	21
Raspberries	
Strawberries	7
Walnuts	14

* PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.
 ** Don't spray only.
 Note: To protect bees, do not spray to fruit trees when trees or adjacent nurseries of weeds around the trees are in bloom.

Crop	Pest	Rate	Timing/Special Directions
Almonds (CA only)	San Jose scale, Pearstone scale	2 tablespoons in 3 gals. of water	Apply only as a dormant spray. Do not apply more than 6 gallons per 600 sq. ft. of orchard. This product may be used on almonds only in CA. Do not use this product on almonds grown in other states.

BEST AVAILABLE COPY

Orchards	Aphids. Aoncho Miteyoutg. Mites (Brown). Two-spined Soderl. Scale Crawlers (Olive, San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 600 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Cane- berries CA, OR, WA only: Black- berries. Raspber- ries. Dew- berries. Logan- berries. Rasp- berries	Aphids. Leafhoppers. Raspberry Fruitworms. Two-spined Soderl. Mites, Thrips	2 tablespoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 6 gallons per 600 sq. ft. of field. Do not make more than 5 applications per season a minimum of 14 days apart. Do not apply within 7 days of harvest. This product may be used on caneberr- ies only in CA, OR and WA. Do not use this product on caneberr- ies grown in other states.
Cherries	Black Cherry Aphids. Cherry Fruitflies. Cherry Fruitworms (Maggot). Fruitree Leafrollers. Mites (Cherry Rust). Scale Crawlers (San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 600 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 7 days apart. Do not apply within 21 days of harvest.
Grapes	Aphids. Leafhoppers. Grape Berry Moths. Grape Leafrollers	4 1/2 teaspoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 3 gallons per 600 sq. ft. of vineyard. Do not make more than 5 applica- tions per season a minimum of 7 days apart. Do not apply within 28 days of harvest.
Decorative	Aphids. Mites (Brown). Two-spined Soderl. Scale Crawlers (Olive, San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 600 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Peaches	Aphids. Mites (Brown). Two-spined Soderl. Scale Crawlers (Olive, San Jose)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 600 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.

Plums and Prunes	Aphids (Leucourt Plum. Mealy Plum, Thistle). Mites (Brown). European Red)	4 teaspoons in 2 gals. of water	Apply as a full cover spray. Do not apply more than 4 gallons per 600 sq. ft. of orchard. Do not make more than 3 applica- tions per season a minimum of 7 days apart. Do not apply within 21 days of harvest.
Straw- berries	Aphids. Two-spined Soderl. Mites. Strawberry Leafrollers	2 tablespoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 3 gallons per 600 sq. ft. of field. Do not make more than 3 applications per season a minimum of 7 days apart. Do not apply within 6 days of harvest.
Walnuts (CA only)	Aphids. Scale Crawlers (Walnut, Italian Pear, Frosted). Spider Mites. Twig Borers	2 tablespoons in 3 gals. of water	Apply as a full cover spray. Do not apply more than 6 gallons per 600 sq. ft. of orchard. Do not make more than 3 applica- tions per season 14 days apart. Do not apply after nuts are open or split. Do not apply within 45 days of harvest. This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states.

BEST AVAILABLE COPY

Vegetables

For foliar or soil applications, D-2-n Lawn and Garden Insect Control should be applied in the volume of water recommended in the table below to assure thorough coverage of the foliage or soil.

Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

Pre-Harvest Interval (PHI)* for Foliar Applications

Crop	PHI* (Days)
Beans ¹ succulent varieties only: Lima, Pole and Snap	7
Broccoli	7
Cabbage	21
Cantaloupes	3
Carrots	14
Cassias	3
Cauliflower	7
Chinese Broccoli, Chinese Cabbage, Collards	10
Corn cobs	3
Cucumbers	7
Enoki (Escarole)	14
Homegrown Melons	3
Kale	10
Lettuce	14
Muskmelons	3
Onions (Bulb and Green), Parsnips	14
Pears ¹	7
Peppers	5
Potatoes (fresh)	35
Radishes, Salsich	14
Squash, Summer	7
Squash, Winter	3
Sweet Corn	7
Tomatoes	1
Turkey (Roots and Tots)	14
Watermelons	3

* PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

¹ Do not use this product on dried pea or bean varieties such as bird beans, dried limas, split peas or dried black-eyed peas.

Vegetables

For application to the foliage, thoroughly spray all plant parts. Application to the soil should be mixed into the soil immediately after application.

Crop	Pest	Rate	Timing/Special Directions
Beans, Lima, Pole and Snap (succulent varieties only)	Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 6 days up to 3 applications per season. Do not apply within 7 days of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms (snap beans only)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-6 inches.

NOTES: (1) Do not apply this product to dried bean varieties such as pinto beans or dried limas. (2) To protect bees, do not apply when bees are in bloom.

Beets, Red (Table)	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.

Broccoli	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 6 applications per season. Do not apply within 7 days of harvest. NOTE: Resistant varieties will not be controlled by this product.
----------	--	--	--

Cabbage	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 6 applications per season. Do not apply within 21 days of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Root Maggots	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-6 inches.

BEST AVAILABLE COPY

Carrots	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 14 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecto- cides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before sowing. Immediately work into soil 2-3 inches for sur- face cutworms, 3-6 inches for subterra- nean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before plant- ing. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-6 inches.
Cauliflower	Aphids. Diamondback Moths. Flea Beetles, imported Cabbageworms. Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 7 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecto- cides.
Chinese Broccoli	Aphids. Diamondback Moths. Flea Beetles. Imported Cabbageworms. Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecto- cides.
Chinese Cabbage (Bok Choy and Napa)	Aphids. Diamondback Moths. Flea Beetles. Imported Cabbageworms. Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecto- cides.
Collards	Aphids. Diamondback Moths. Flea Beetles. Imported Cabbageworms. Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecto- cides.

Corn (see sweet corn)	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 7 days of harvest.
Cucumbers	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 7 days of harvest.
Endive (Escarole)	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 6 applica- tions per season. Do not apply within 14 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecto- cides.
Kale	Aphids. Diamondback Moths. Flea Beetles. Imported Cabbageworms. Dipterous Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 10 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecto- cides.
Lettuce	Aphids. Flea Beetles. Leafhoppers. Dipterous Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 14 days of harvest. NOTE: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecto- cides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before sowing. Immediately work into soil 2-3 inches for sur- face cutworms, 3-6 inches for subterra- nean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before sowing. Immediately work into soil 4-6 inches.
Melons (Cantaloupe, Cassia, Cren- shaws, Honey- dews, Musk- melons, Persians, and Hybrids of these) and Water- melons	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applica- tions per season. Do not apply within 3 days of harvest.

BEST AVAILABLE COPY

Onions (Bulb and Green)	Onion Thrips	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not spray within 14 days of harvest.
Peas (succulent only)	Aphids. Cucumber Beetles. Grasshoppers. Leafhoppers. Mexican Bean Beetles. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Allow at least 7 days between the last application and harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-6 inches.
NOTES: (1) Do not apply this product to dried pea varieties such as split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if bees or weeds in the treatment area are in bloom.			
Peppers	Aphids. Colorado Potato Beetles. Flea Beetles. Leafhoppers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not spray within 5 days of harvest.
Potatoes	Aphids. Flea Beetles. Leafhoppers. Colorado Potato Beetles. Southern Armyworms. Diocterus Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not spray within 35 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

Radishes	Aphids. Flea Beetles. Leafhoppers. Imported Cabbageworms. Diocterus Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 3 applications per season. Do not spray within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
Squash	Aphids. Flea Beetles. Leafhoppers. Diocterus Leafminers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 6 applications per season. Do not spray within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Squash, Summer	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 6 applications per season. Do not spray within 7 days of harvest.
Squash, Winter	Aphids. Thrips. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 6 applications per season. Do not spray within 3 days of harvest.
Sweet Corn	Aphids, Thrips. Flea Beetles. Grasshoppers. Spider Mites	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Corn may be picked 7 days following last application.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-6 inches.
NOTE: To protect bees, do not apply to corn foliage during the pollen shed period.			

BEST AVAILABLE COPY

Tomatoes	Aphids. Colorado Potato Beetles. Fall Armyworms. Flea Beetles. Leafhoppers. Southern Armyworms. Vinegar Flies Drosophila spp.	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 1 day of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 4-6 inches.
Turnips and Cauliflowers	Aphids. Flea Beetles. Leafhoppers. Imported Cabbageworms. Dorset Leafminers	2 tablespoons in 3 gals. of water to cover 1,000 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. NOTE: Resistant leafminers will not be controlled by diazinon or other type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 1/2 gals. of water	Broadcast to the soil just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.

Storage and Disposal

Storage
Do not store near heat or open flame. Store in original container in an area inaccessible to children and pets. **USE CONTENTS WITHIN 6 MONTHS AFTER OPENING.**

Disposal
Do not reuse empty container. Thoroughly rinse container, securely wrap it in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

Precautionary Statements

Hazards to Humans and Domestic Animals
CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid breathing spray mist. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils such as spoons and measuring cups must not be used for food purposes after use in measuring pesticides.

Statement of Practical Treatment
If swallowed: Call a physician immediately. **DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE.** Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled: Remove victim to fresh air and apply artificial respiration if needed. Get medical attention.

If on skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If in eyes: Rinse eyes with plenty of water and call a physician immediately.

Notes to Physician: This product is an organophosphorus insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is essential. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains aromatic petroleum hydrocarbons which may present an aspiration hazard. Gastric lavage may be indicated if product was taken recently.

Environmental Hazards
This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lands in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. A.C.Z. overloading of sprays. For lawns, where irrigation (watering) is recommended in the direction of waterfowl, i.e., duck or geese, can be expected in the treat. (A.Z. after treatment (except spot treatment), water lawn with at least 1/2 inch of water immediately after spraying the product. However, stop watering before quacking occurs. Keep A.Z. of lakes, streams, ponds, total manure and excreta. Do not apply directly to water.

Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on the label. Do not apply where fish, shrimp, crab, and other aquatic life are over-valued resources. Do not contaminate water by cleaning or equipment or disposal of equipment wash water.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to be applied to crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards
Do not use, pour, spill, or store near heat or open flame.

BEST AVAILABLE COPY