

DIRECTIONS FOR USE

FOR DIRECT APPLICATION TO DAIRY CATTLE AND BEEF CATTLE, FOR CONTROL OF HORNFLIES: Apply 2 heaping tablespoonsful (2 oz.) to the backs and necks of infested animals. Rub lightly into hair. Repeat application in 10 to 14 days if necessary. On dairy animals, repeat no more often than once in every three week period, if necessary.

FOR CONTROL OF LICE AND TICKS ON BEEF CATTLE: Treat entire animal very thoroughly. Repeat treatment at two week intervals if necessary. May be used on nonlactating dairy animals for lice control up to within two weeks of freshening.

AS AN AID FOR THE CONTROL OF FACE FLIES ON BEEF CATTLE: Dust the faces of cattle thoroughly. Frequent repeated applications may be necessary.

FOR CONTROL OF HOG LICE: Apply to hogs, pens, and litter. One treatment may be sufficient. Treat animals thoroughly. Repeat in 10 days or thereafter as needed.

LICE AND KEDS ON SHEEP: Treat animals thoroughly, preferably following shearing. Repeat as necessary.

CAUTION: Avoid prolonged breathing of dust. Avoid contact with skin; wash thoroughly after using. Keep out of reach of children. Do not contaminate feed, water, or foodstuffs. Do not apply within five hours before, or during milking. Do not treat animals under one month of age.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

1023-7112

1973

70-183

KILL-KO

MAL-THOX

H D C
HORSES DAIRY CATTLE

DUST

LIVESTOCK

ACTIVE INGREDIENTS

Malathion (0.6 g/methoxyphenyl)
Methoxyphenyl (0.6 g/methoxyphenyl)
Methoxyphenyl (0.6 g/methoxyphenyl)

INERT INGREDIENTS

*Equivalent to 4.4% 2,2 bis (p Methoxyphenyl)
1,1,1 trichloroethane and 0.6% related products.

KEEP OUT OF REACH OF CHILDREN.
SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

NET WEIGHT ONE POUND

