

DIRECTIONS FOR USE

Apply 1/2 lb. of ZECTRAN Snail & Slug Killer Meal at the rate of 1/2 lb. per 100 sq. ft. of infested area. Water the infested area thoroughly before application. Then spread the meal evenly over the area where snails travel and congregate. Special care should be taken to treat bedding plants or other low growing plants where pests feed and breed. Pests stop feeding soon after treatment but kill may be delayed for 3 to 5 days. Re-treat every 3 weeks or as needed.

CAUTION

This product is toxic to fish, wildlife and birds. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Bury waste in non-cropland away from water supplies. Dispose of empty container by burning or by burying with waste. Apply this product only as specified on this label.

Wash thoroughly after handling. Avoid contamination of food, food-stuff and domestic water supplies. If swallowed, induce vomiting by giving an emetic, such as two tablespoonful of salt in a glass of water. Call a physician immediately.

NOTE TO PHYSICIAN: Active ingredient has cholinergic effects. Treat symptomatically. Atropine is an antidote.

For use on lawns, flower gardens, other ornamental plantings and on ground covers including grass and Dichondra lawns.



ZECTRAN*

SNAIL AND SLUG KILLER MEAL

Kills Sowbugs
(Pillbugs) Too



FOR USE IN FLOWER GARDENS, OTHER ORNAMENTAL
PLANTINGS AND ON GROUND COVERS INCLUDING GRASS
AND DICHONDRA LAWNS.

ACTIVE INGREDIENT

4-(dimethylamino)-3, 5-xylol methylcarbamate

INERT INGREDIENTS

*Trademark of The Dow Chemical Company

See also the back of each of our
ZECTRAN products for more information.

DIRECTIONS FOR USE

Apply KILL-KO ZECTRAN Snail & Slug Killer Meal at the rate of 12 ounces per 900 square feet of treatment area. Water the infested area thoroughly before application, then spread the meal evenly over the area where pests travel and congregate. Special care should be taken to treat bedding plants or other low growing plants where pests feed and hide. Pests stop feeding soon after treatment but kill may be delayed for 3 to 5 days. Re-treat every 3 weeks or as needed.

CAUTION

This product is toxic to fish, wildlife and birds. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Bury waste in non-cropland away from water supplies. Dispose of empty container by burning or by burying with waste. Apply this product only as specified on this label.

Wash thoroughly after handling. Avoid contamination of food, food-stuff and domestic water supplies. If swallowed, induce vomiting by giving an emetic such as two tablespoonsful of salt in a glass of warm water. Call a physician immediately.

NOTE TO PHYSICIAN: Active ingredient has cholinergic effects. Treat symptomatically. Atropine is an antidote.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

CAUTION

KEEP OUT OF REACH OF CHILDREN.

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing.

2455 714

U.S. Patent No. 3,084,098
EPA REG. NO. 70-189



ZECTRAN*

SNAIL AND SLUG KILLER MEAL

Kills Sowbugs
(Pillbugs) Too



FOR USE IN FLOWER GARDENS, OTHER ORNAMENTAL
PLANTINGS AND ON GROUND COVERS INCLUDING GRASS
AND DICHONDRA LAWNS.

ACTIVE INGREDIENT

4-(dimethylamino)-3, 5-xylyl methylcarbamate

INERT INGREDIENTS

Trade Mark of the Dow Chemical Company

CAUTION KEEP OUT OF REACH OF CHILDREN
SEE OTHER PRECAUTIONS ON SIDE PANEL

NET WEIGHT 12 OUNCES

DOW COMPANY
DOWNEY, CALIFORNIA 90241

DOW COMPANY
DOWNEY, CALIFORNIA 90241