

# RIGO RAT BAIT-F

ACTIVE INGREDIENTS:  
3-(alpha-acetylfurfuryl)-4-Hydroxycoumarin 0.025%  
INERT INGREDIENTS: 99.975%

**CAUTION:**  
**KEEP OUT OF REACH OF CHILDREN**

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

In case bait is eaten by humans, call physician at once. Treat as for overdose of Dicoumarol.

For Repackaging Only By Firms Holding An Approved E.P.A.  
Registration.  
Not For Resale Under This Label.

EPA REG. NO. 70-187

MANUFACTURED BY

**Rigo Company**  
NASHVILLE, TENN. 37202

NET WEIGHT . . . POUNDS



**KILL-KO**

JAN 28 1974  
**20% LINDANE**

**(Emulsifiable Concentrate)**

**CAUTION**

Keep out of reach of children. Harmful if swallowed. For first aid in case any of the insecticide is swallowed, give a tablespoonful of salt in a glass of warm water. Repeat until vomit fluid is clear. Keep patient prone and warm. Call a physician immediately. There may be an aspiration hazard associated with emesis.

Avoid inhalation of spray mist. Avoid skin contact. Harmful vapors may be absorbed by the skin. Wash thoroughly with soap and water. For the eyes, get medical attention; flush with water for 15 minutes.

Avoid contamination of feed or food-stuffs. Food utensils such as teaspoons or measuring cup should not be used for food purposes after using with insecticide.

This product is toxic to fish and wildlife. B. feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of waste.

Destroy this container when empty. Do not reuse.

**DO NOT USE, POUR, SPILL, SPRAY OR STORE NEAR HEAT OR OPEN FLAME**

**HOW KILL-KO LINDANE EMULSIFIABLE CONCENTRATE KILLS INSECTS**

Kill-Ko 20% Lindane Concentrate is effective because it kills by two methods, "Contact" and "Stomach Poisoning". This quality is a help in killing insects that tend to bite or burrow into plants.

**DIRECTIONS FOR USE**

Note Caution Statement and read directions thoroughly before using. Apply the product only as specified on the label. Before using KILL-KO 20% LINDANE EMULSIFIABLE CONCENTRATE on livestock animal, it is recommended that you consult your State Agricultural Extension Service for use, the instructions on page 1 of the label and other information.

**ANIMAL USES**

**SHEEP & SHEEP LAMB** - For control of lice, ticks, and mites on sheep and lambs, use Dilution No. 1. For control of ticks on lambs, use Dilution No. 2. For control of ticks on sheep, use Dilution No. 3. For control of ticks on lambs, use Dilution No. 2. For control of ticks on sheep, use Dilution No. 3.

particularly desirable for sheep.

**DOGS:** For sarcoptic mange apply Dilution "1" once a week for two or more treatments as needed. For fleas, ticks, or lice, use Dilution "A" and apply at 7 to 10 day intervals as needed but never more often than once per week. For kennels spray walls thoroughly with Dilution No. 3.

**CAUTION:** Do not use on cats or household pets other than dogs. Do not use on nursing puppies or bitches with nursing pups. Never use Dilution No. 3 directly on the animals.

**HOGS:** For mange (Sarcoptic), lice and ticks spray thoroughly with Dilution A. Repeat application at weekly intervals for two or more treatments as needed.

**BEEF CATTLE & HORSES:** For ticks, lice and horn fly use Dilution No. 2. Apply to adult animals at the rate of about two gallons of diluted spray per head with only two sprays, a week apart.

**RESIDUAL SPRAY AND PAINT**

**RESIDUAL SPRAY:** For use in barns, calf pens and manure piles, in the control of flies, stable flies, roaches, ants and mosquitoes, treat walls, cracks and crevices with Dilution No. 3.

**RESIDUAL PAINT:** Dilute one part KILL-KO LINDANE EMULSIFIABLE CONCENTRATE with nine parts of water and use as a paint. Paint windows, doors and other places where flies congregate.

**CAUTION:** Do not use in poultry houses, dairy barns, milk rooms or on dairy cattle or poultry. Do not use this product as a dip or in back rubbers. Do not apply with lime or fresh washdown, or with other alkaline material. Treatment may be advisable to prevent infestation of the herd. In such case, the amount per animal should be no more than a quart and proportionately less for small animals. Do not treat animals under three months of age. Do not treat cows within 2 weeks of parturition or at least 2 weeks thereafter. Avoid treating animals in cold, rainy weather. Do not exceed Dilution "A" on young animals such as calves. Avoid treating overfatted or sick animals. Do not spray more than 30 days after parturition.

**FRUIT TREES**

For control of fruit tree pests, use Dilution No. 1. For control of fruit tree pests, use Dilution No. 1. For control of fruit tree pests, use Dilution No. 1. For control of fruit tree pests, use Dilution No. 1. For control of fruit tree pests, use Dilution No. 1.