


KILL-KO



protects your fruit trees

CONTAINS

Methoxychlor Malathion Captan

An insecticide and fungicide for insect and disease control on the fruit trees

ACTIVE INGREDIENTS:

Technical Methoxychlor*	10.0%
Captan (N-trichloromethylthio-4-cyclohexene-1, 2-dicarboximide)	7.0%
Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate)	5.0%
INERT INGREDIENTS:	78.0%
	100.0%

*Equivalent to 8.8% 2,2-bis(p-methoxyphenyl)-1, 1, 1-trichloroethane and 1.2% other isomers and related compounds.

Out of the Reach of Children

NET WEIGHT 12 OUNCES

Manufactured By
RIGO COMPANY
BUCKNER, KY. 40010

DIRECTIONS

BEFORE USE READ CAUTION AND NOTICE STATEMENTS CAREFULLY.

SPRAY APPLICATION: 12 oz. of this material will make 7½ gallons of fruit tree spray. For smaller quantities, use 5 level tablespoonfuls to each gallon of water. Mix well. Agitate solution while spraying. Apply spray to trees thoroughly.

APPLES, PEARS: To control Bitter Rot with Scab and Brooks Fruit Spot, Woolly and Green Aphids, Pacific, Clover and Red Spider Mites, Codling Moth, Forbes Scale, Plum Curculio, Apple Maggot, Japanese Beetle, Red Banded Leaf Roller and Bitter Rot, apply as follows:

1. When blossom stems separate and before blossoms open.
2. When 75% of the petals have fallen.
3. 10 days later.
4. 10 days to 2 weeks later.
5. Continue at 2-week intervals to 21 days before harvest.

Do not use on D'Anjou pears. Russetting may be produced on Bosc pears. Red delicious and other sensitive varieties may be injured by early season application.

PEACHES: To control Brown Rot, Peach Scab, European Red Mite, Red Spider, Plum Curculio, Oriental Fruit Moth, Japanese Beetle, European Fruit Scale, Pacific and Clover Mites, apply as follows:

1. Full bloom.
2. When shucks begin to split.
3. 10 days later.
4. 10 days later.
5. 2 weeks later.
6. Do not apply within 21 days of harvest.

PLUMS (PRUNES): To control Pacific, European and Red Spider Mites, Plum Curculio, Mealy Plum and Green Peach Aphids and Japanese Beetle, apply as follows:

1. When trees are in full bloom.
2. When shucks are falling.
3. 10 days later.
4. 2 weeks later.
5. Continue at 2-week intervals if necessary. Do not apply later than 21 days before harvest.

If control of disease on above fruit crops is necessary within 21 days of harvest, apply a wettable sulfur spray. Remove residues by washing, wiping, brushing or other effective means.

CAUTION

Harmful if swallowed. Avoid inhalation of dust or spray mist. Avoid contact with skin; wash skin and hands thoroughly after using. Do not get in eyes. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residues in excess of official tolerances.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Education Service.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

NOTICE: This makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

343



ACCEPTED

APR 2 1974

UNDER THE
 FUNCTIONS OF THE
 FOR ECONOMIC POLICY
 ED UNDER NO 70-190 SUBJECT
 TO ATTACHED COMMENTS.