


70-190 14112

103
0



K I L L - K O
FRUIT TREE SPRAY

An Insecticide and Fungicide for Insect
and Disease Control on Fruit Trees

Contains Methoxychlor*, Malathion, Captan

ACTIVE INGREDIENTS:

- Technical Methoxychlor* 10.0%
- Captan (N-trichloromethylthio-4-cyclohexene-1; 2-dicarboximide) 7.0%
- Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate). 5.0%

INERT INGREDIENTS78.0%
 100.0%

*Equivalent to 8.8% 2,2-bis(p-methoxyphenyl)-1, 1, 1-trichloroethane and 1.2% other isomers and related compounds.

CAUTION: (14 pt.)


KEEP OUT OF REACH OF CHILDREN (10 pt.)

EPA REG. NO. 70-190
EPA EST. NO. 70-KY-1

NET WEIGHT
____ OUNCES

See Back Panel for Other Precautionary Statements

Manufactured by
RIGO COMPANY
Buckner, Kentucky 40010



503

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Avoid inhalation of dust or spray mist. Avoid contact with skin; wash skin and hands thoroughly after using. Do not get in eyes. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residues in excess of official tolerances.

Environmental Hazards

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Disposal (10 pt.)

Do not reuse empty container. Wrap container and place in trash.


DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SPRAY APPLICATION: 10 oz. of this material will make 6 1/2 gallons of fruit tree spray. For smaller quantities, use 5 level tablespoonfuls to each gallon of water. Mix well. Agitate solution while spraying. Apply spray to trees thoroughly.

APPLES, PEARS: To control Bitter Rot, Scab, Brooks Fruit Spot, Woolly and Green Aphids, Pacific, Clover and Red Spider Mites, Codling Moth, Forbes Scale, Plum Curculio, Apple Maggot, Japanese Beetle, and Red Banded Leaf Roller, apply as follows:

1. When blossom stems separate and before blossoms open.
2. When 75% of the petals have fallen.
3. 10 days later.
4. 10 days to 2 weeks later.
5. Continue at 2-week intervals to 21 days before harvest.



KILL-KO FRUIT TREE SPRAY

Page 3

Do not use on D'Anjou pears. Russetting may be produced on Bosc pears. Red delicious and other sensitive varieties may be injured by early season application.

PEACHES: To control Brown Rot, Peach Scab, European Red Mite, Red Spider, Plum Curculio, Oriental Fruit Moth, Japanese Beetle, European Fruit Scale, Pacific and Clover Mites, apply as follows:

1. Full bloom.
2. When shucks begin to split.
3. 10 days later.
4. 10 days later.
5. 2 weeks later.
6. Do not apply within 21 days of harvest.

PLUMS (PRUNES): To control Pacific, European and Red Spider Mites, Plum Curculio, Mealy Plum and Green Peach Aphids and Japanese Beetle, apply as follows:

1. When trees are in full bloom.
2. When shucks are falling.
3. 10 days later.
4. 2 weeks later.
5. Continue at 2-week intervals if necessary. Do not apply later than 21 days before harvest.

If control of disease on above fruit crops is necessary within 21 days of harvest, apply a wettable sulfur spray. Remove residues by washing, wiping, brushing or other effective means.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.