

**ACCEPTED**  
 APR 9 1976  
 Under the Federal Insecticide,  
 Fungicide, and Rodenticide Act,  
 as amended, for the pesticide  
 registered under  
 EPA Reg. No. 70-190



# 3% Rabon® Dust

## A Ready-To-Use Insecticide Dust FOR LIVESTOCK AND POULTRY

**ACTIVE INGREDIENT**  
 2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate ..... 3.00%  
**INERT INGREDIENT** ..... 97.00%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:**

**STATEMENT OF PRACTICAL TREATMENT**

If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician.

**PRECAUTIONARY  
 STATEMENTS**

**HAZARD TO HUMANS:**  
 May be fatal if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings.

**ENVIRONMENTAL HAZARDS!**  
 This product is toxic to fish. Keep out of lakes.

**FOR POULTRY:**

Controls northern fowl mites, chicken mites, and lice.

TYPE HOUSING	APPLICATION	REMARKS
WIRE CAGES	1 lb/300 birds  Plunger, rotary type duster or shaker-can-duster.	For individual treatment direct dust to vent and fluff area. Group treatment may be preferred. Dust should reach skin. Do not repeat more often than every 14 days.  Wire rungs and corners should also be treated.
<b>NOTE:</b> For northern fowl mites on roosters, thorough individual application of the dust will assure long-lasting control and reduce reinfestation of breeding flocks.		
FLOOR		

2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate ..... 3.00%  
 INERT INGREDIENT ..... 97.00%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:**

**STATEMENT OF PRACTICAL TREATMENT**

If the material gets into the eyes, wash with plenty of water for 15 minutes; if irritation persists, see a physician.

**PRECAUTIONARY STATEMENTS**

**HAZARD TO HUMANS:**

May be fatal if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat and open flame. Do not apply in dwellings.

**ENVIRONMENTAL HAZARDS!**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**USE DIRECTIONS**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. **PESTICIDE DISPOSAL:** Excess pesticide that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. **CONTAINER DISPOSAL:** Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

6333-7105  
 Registered Shell Trademark  
 EPA Est. 70-KY-1  
 EPA Reg. No. 70-192-11885

**NET CONTENTS 50 LBS.**

**FOR POULTRY:**

Controls northern fowl mites, chicken mites, and lice.

TYPE HOUSING	APPLICATION	REMARKS
<b>WIRE CAGES</b>	1 lb/300 birds  Plunger, rotary type duster or shaker-can-duster.	For individual treatment direct dust to vent and fluff area. Group treatment may be preferred. Dust should reach skin. Do not repeat more often than every 14 days.  Wire rungs and corners should also be treated.
<b>NOTE:</b> For northern fowl mites on roosters, thorough individual application of the dust will assure long-lasting control and reduce reinfestation of breeding flocks.		
<b>FLOOR MANAGEMENT</b>		
Litter	1 lb/100 sq. ft. Plunger or rotary type duster.	Treat evenly and thoroughly. Also treat roosts, cracks and crevices where pests may hide.
Dust Box	2 lb/100 birds	Use box about 2 ft. by 2 ft. by 1 ft. deep.
Roost Paint	1 lb/100 ft. make thick slurry by mixing 1 lb. of dust with 1 pint of water continually stir.	Treat roosts thoroughly, particularly the cracks and crevices.

**FOR BEEF CATTLE & DAIRY CATTLE:**

To control HORN FLIES and LICE and an aid in the control of FACE FLIES. Apply by hand or through the use of self-treating dust bags. There is no withholding period from last application to slaughter. For Horn Flies and Face Flies on beef and dairy cattle, use as follows:

**HAND DUSTING** — Apply approximately 2 oz. of dust by shake can, rotary duster or by spoon to the upper portions of the back, neck and poll, and to the face as an aid in the control of face flies. Rub in lightly to carry the dust beneath the hair. Repeat as necessary.

**SELF-TREATING DUST BAG —**

**FORCED USE** — Puncture the paper sleeve and insert into the dusting bag. Hang in barn door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or burlap bags from weather.

**FREE CHOICE USE** — Use the same dust bags as above but place in loafing sheds, holding pens, feedlots, near watering holes or other areas where cattle gather. The Free Choice use aids in the control of Lice.

**FOR SWINE: To control lice.**

**HAND-DUSTING** — Apply 3-4 oz. of dust by conventional hand or powder duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and bedding may be treated. One lb. of 3% Dust should be applied per 150 square feet of bedding.

Manufactured By  
**RIGO COMPANY**  
 BUCKNER, KENTUCKY 40010