



KO

KO

### HUMAN CAUTION

HARMFUL IF INHALED OR SWALLOWED DO NOT breathe dust. Avoid contact with skin. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT store near feed or foodstuffs.

### ENVIRONMENTAL CAUTION

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

### NOTICE

Seller makes no warranty, expressed or implied, concerning use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

### GENERAL INFORMATION

KILL KO 1 1/4% SEVIN DUST controls a wide range of insect pests. At first signs of insect infestation, preferably before damage from feeding occurs, apply thoroughly to upper and lower leaf surfaces. If needed, repeat applications every 7-10 days. DO NOT use with lime or other alkaline material. May be applied up to harvest except where noted.

### DIRECTIONS

BEANS -- For control of Mexican bean beetle, leafhopper, flea beetles, apply approximately 1 1/2 lbs./1000 sq. ft. DO NOT feed treated bean vines to dairy or meat animals.

LETTUCE -- For control of lygus and stick bugs, apply approximately 2 to 2 1/2 lbs./1000 sq. ft. DO NOT apply to head lettuce within 3 days of harvest or within 14 days to leaf lettuce.

CORN -- Use approximately 2 to 2 1/2 lbs. per 1,000 sq. ft. for corn earworm, fall armyworm, flea beetles, Japanese Beetles, Sap Beetles, leafoppers and European corn borer. For corn borer and larvae in whorls (budworm damage) and for foliage feeders, make one or more applications to entire plant beginning when borer damage threatens depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experimental station for proper timing of treatments. For home or small garden use, apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until control is achieved. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Re-apply immediately after rain. Allow 7 days before harvest or grazing of forage. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS -- For control of Colorado potato beetle, leafhoppers and flea beetles, apply approximately 1/4 to 1 1/4 lbs./1000 sq. ft. For control of tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, stink bug and lygus bugs, apply approximately 2 to 2 1/2 lbs./1000 sq. ft.

CUCUMBERS, SUMMER SQUASH -- For control of pickleworm, melonworm, cucumber beetles, squash bugs, flea beetle, apply approximately 1 1/4 lbs./1000 sq. ft.

GRAPES -- For control of leafhopper, Japanese beetle, grape berry moth, apply approximately 2 1/4 lbs./1000 sq. ft.

STRAWBERRIES -- For control of meadow spittlebug, leaf roller apply approximately 1 1/2 lbs./1000 sq. ft. DO NOT apply within 1 day of harvest.

ROSES, AZALEAS, SHRUBS AND OTHER ORNAMENTALS -- For control of Japanese beetles,

