

6/20/97

PM 19

70-244

pg 1/4

RIGO 5% SEVIN® BAIT

For Control of Listed Insect Pests

Active Ingredient:

Carbaryl ((1-Naphthyl N-methylcarbamate) 5%

Inert Ingredients: 95%

Total: 100%

Sevin® is a registered trademark of Rhone-Poulenc for carbaryl insecticide

EPA Reg. No. 70-244

EPA Est. No. 769-GA-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Side panel For Additional Precautionary Statements

STATEMENT OF PRACTICAL TREATMENT

If On Skin: Wash thoroughly with soap and water.

If In Eyes: Flush with water for at least 15 minutes and get medical attention.

If Swallowed: Induce vomiting by drinking 1 or 2 glasses of water and touching back of throat with finger or blunt object. Call a physician or Poison Control Center immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

Note To Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase-inhibiting drugs.

Net Contents: _____

Manufactured By:
Rigo Company
Louisville, KY 40222

| |
|---|
| ACCEPTED |
| JUN 20 1997 |
| Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 70-244 |

PRECAUTIONARY STATEMENTS
Hazards To Humans and Domestic Animals
CAUTION

Harmful if swallowed or inhaled. Avoid breathing of dust. Do not take internally. Avoid contact with skin. Avoid contamination of food and feed.

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until dust has settled.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic organisms. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use when value of bees as pollinators is more important than insect control. Before application, warn beekeepers to locate hives beyond bee flight range until 1 week after application or to take other equally effective precautions.

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable SEVIN residues when SEVIN is used in accordance with label directions. If SEVIN insecticide is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in residue in milk or meat in excess of established tolerances.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Broadcast application treatments may be repeated once a week. Dosages of bait from 1/2 to 1 pound per 1,000 sq. ft. will control cutworms, armyworms, field crickets, darkling beetles, grasshoppers, sowbugs and ants.

NOTE: To avoid possible injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

Wearing household latex or rubber gloves, use a shaker can, or rotary type applicator. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow any dust to settle in treated areas before reentering. For plants above chest height, consider using a liquid carbaryl product. Do not repeat application more than once a week.

| Crop | Dosage Rate Pounds/1000 sq. ft. | Days Between Final Application and Harvest |
|--|------------------------------------|---|
| Cucumber, Squash, Melons, Peanuts | 1/2 | None |
| Peas | 1/4 | None |
| Beans, Carrots, Eggplants, Okra, Peppers, Potatoes, Tomatoes | 1 | None |
| Corn and Sweet Corn | 1/2 - 1 | None |
| Asparagus, Strawberries | 1 | 1 Day |
| Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Head Lettuce, Garden Beets (roots), Horseradish, Parsnips, Radishes, Rutabagas, Turnips (roots) | 1 | 3 Days |
| Collards, Endive (escarole), Garden Beets (tops), Kale, Leaf Lettuce, Parsley, Spinach, Swiss Chard, Turnips (tops) | 1 | 14 Days |
| Dandelions | 1 | 14 days |

| Crop | Dosage Rate Pounds/1000 sq. ft. | Days Between Final Application and Harvest |
|--|------------------------------------|---|
| All insects listed above (except darkling beetles) for the following crops: Apples, Cherries, Peaches, Pears, Plums To avoid possible injury to McIntosh and York apples, do not apply before the 2nd cover. To avoid thinning apples, delay use at least 30 days following full bloom | 1 | 1 Day |
| Apricots, Nectarines | 1 | 3 Days |
| Citrus | 1 | 5 Days |
| Grapes | 1 | None |
| Annual plants, herbaceous plants and shrubs, lawns or turf, flower and herb gardens, ornamental trees, shrubs, and flowers, and soil surfaces around the outside of buildings (including houses) | 1 | None |

STORAGE AND DISPOSAL

Storage: Store in original container in a cool, dry place out of reach of children and pets.

Disposal: Do not reuse empty container. Wrap empty container in newspaper and put in trash.

Notice: Seller makes no warranty, expressed or implied, concerning the use of this product other than stated on the label. Buyer assumes all risk of use and/or handling of the material when such use and/or handling is contrary to label instructions.

05/97